

Applicants: Zhongyi Li et al.

Serial No.: 10/577,564

Filed: April 26, 2006

Exhibit 5

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 May 2002 (16.05.2002)

PCT

(10) International Publication Number
WO 02/37955 A1

(51) International Patent Classification⁷: **A01H 5/00**,
5/10, C12N 15/29, C08L 3/02

BATEY, Ian, Leslie [AU/AU]; CSIRO Australia, Division
of Plant Industry, Cnr Clunies Ross Street and Grain Drive,
Canberra, ACT 2601 (AU).

(21) International Application Number: PCT/AU01/01452

(22) International Filing Date:
9 November 2001 (09.11.2001)

(74) Agent: **A.P.T. PATENT AND TRADE MARK ATTOR-
NEYS**; GPO Box 772, Adelaide, S.A. 5001 (AU).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PR 1371 9 November 2000 (09.11.2000) AU
PR 1370 9 November 2000 (09.11.2000) AU
PR 1372 9 November 2000 (09.11.2000) AU
PR 1373 9 November 2000 (09.11.2000) AU

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN,
YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).

(71) Applicant (*for all designated States except US*): **COM-
MONWEALTH SCIENTIFIC AND INDUSTRIAL
RESEARCH ORGANISATION** [AU/AU]; Limestone
Avenue, Campbell, ACT 2612 (AU).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **MORELL,
Matthew, Kennedy** [AU/AU]; CSIRO Australia, Di-
vision of Plant Industry, Cnr Clunies Ross Street and Grain
Drive, Canberra, ACT 2601 (AU). **TOPPING, David**
[AU/AU]; CSIRO Australia, Division of Health Sciences
and Nutrition, Kintore Avenue, Adelaide, S.A. 5000 (AU).

Published:

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: **BARLEY WITH REDUCED SSII ACTIVITY AND STARCH CONTAINING PRODUCTS WITH A REDUCED AMY-
LOPECTIN CONTENT**

(57) Abstract: Barley with reduced SSII activity has a starch structure with reduced amylopectin content and a consequent high relative amylose content. Additionally the grain has can have a relatively high β glucan content. The structure of the starch may also be altered in a number of ways which can be characterised by having a low gelatinisation temperature but with reduced swelling. The viscosity of gelatinised starch of the starch is also reduced. There is a chain length distribution characterised by a low crystallinity of the starch. The starch is also characterised by having high levels of V form starch crystallinity. The dietary fibre content of the starch is high. This

Applicants: **Zhongyi Li et al.**
Serial No.: 10/577,564
Filed: April 26, 2006
Exhibit 5

WO 02/37955 A1

BARLEY WITH REDUCED SSII ACTIVITY AND STARCH AND STARCH CONTAINING PRODUCTS WITH A REDUCED AMYLOPECTIN CONTENT.

This invention relates to a barley plant with a reduced SSII enzyme activity leading to a starch
5 having reduced amylopectin content. The invention also relates to starch and grain and food
products obtained therefrom.

BACKGROUND OF THE INVENTION

One finding in nutritional science is that resistant starch has important implications for bowel
10 health, in particular health of the large bowel. The beneficial effects of resistant starch result
from the provision of a nutrient to the large bowel wherein the intestinal microflora are given
an energy source which is fermented to form *inter alia* short chain fatty acids. These short
chain fatty acids provide nutrients for the colonocytes, enhance the uptake of certain nutrients
across the large bowel and promote physiological activity of the colon. Generally if resistant
15 starches or other dietary fibre is not provided the colon is metabolically relatively inactive.

There has in recent years been a direction to look at providing for resistant starches from
various sources to address bowel health. Accordingly high amylose starches have been
developed in certain grains such as maize for use in foods as a means of promoting bowel
20 health.

The physical structure of starch can have an important impact on the nutritional and handling
properties of starch for food products. Certain characteristics can be taken as an indication of
starch structure including the distribution of amylopectin chain length, the degree of
25 crystallinity and the presence of forms of crystallinity such as the V-complex form of starch
crystallinity. Forms of these characteristics can also be taken as indicator of nutritional or
handling properties of foods containing these starches. Thus short amylopectin chain length
may be an indicator of low crystallinity and low gelatinisation and is also thought to have a
correlation with reduced retrogradation of amylopectin. Additionally shorter amylopectin chain
30 length distribution is thought to reflect organoleptic properties of food in which the starch is
included in significant amounts. Reduced crystallinity of a starch may also be indicative of a
reduced gelatinisation temperature of starch and additionally it is thought to be associated with
enhanced organoleptic properties. The presence of V-complex crystallinity or other starch
associated lipid will enhance the level of resistant starch and thus dietary fibre.

of those foods or on the functional characteristics of those components in the preparation or structure of the foods.

5 Whilst modified starches or β glucans, for example, can be utilised in foods that provide functionality not normally afforded by unmodified sources, such processing has a tendency to either alter other components of value or carry the perception of being undesirable due to processes involved in modification. Therefore it is preferable to provide sources of constituents that can be used in unmodified form in foods.

10 The barley variety MK6827 is available from the Barley Germplasma Collection (USDA-ARS National Small Grain Germplasma Research Facility Aberdeen, Idaho 831290 USA). The grain of MK6827 is shrunken and has a highly coloured husk and an elongate shape and, in the hands of the inventors, this grain is very difficult to process including being very resistant to milling. The properties of MK6827 grain had not been characterised before, nor had the
15 nature of the mutation been ascertained nor is it considered suitable for producing food.

SUMMARY OF THE INVENTION

This invention arises from the isolation and characterisation of SSII mutant of barley plants the grain of which is found to contain starch that has reduced amylopectin content and therefore
20 high relative levels of amylose and therefore has elevated levels of dietary fibre.

The grain of the mutant and grain from crosses into certain genetic backgrounds additionally has an elevated level of β glucan. The combination of elevated β glucan level and resistant starch contributing to high dietary fibre is thought by the inventors to be unique to the present
25 invention.

Additionally, at least in some genetic backgrounds, it is found that grain from such mutants contain starch that have high relative levels of amylose, and also have low gelatinisation temperatures. The low swelling characteristics of such starch during and following
30 gelatinisation also has advantages in certain dietary and food processing applications.

Furthermore, grain from such mutants are found to contain starch that have high relative levels of amylose, the amylose levels found are higher than 50% of the starch content which is a level never before found in unmodified starch derived from barley.

by (+) yielded the Himalaya PCR pattern and lines denoted by (○) gave the 292 PCR result. Panel (A), the seed length to thickness ratio plotted against the percentage of starch chains with DP between 6 and 11; Panel (B) seed weight plotted against the percentage of starch chains with DP between 6 and 11

Figure 9 Sequence of a barley SSII cDNA (SEQ ID NO 1) from the cultivar Himalaya

Figure 10 The structure of the SSII genes from (1) *T. tauschii* (diploid wheat), (2) barley cultivar Morex. The thick lines represent exons and the thin lines introns. The straight line underneath each example indicates the region of the gene sequences. The dotted line represents a region of the barley SSII gene, from intron 7, that has not been sequenced but has been determined by PCR analysis to be approximately 3 kb in length.

Figure 11 Comparisons of the predicted SSII cDNAs from MK6827 (SEQ ID NO 2), Morex (SEQ ID NO 3) and 292 (SEQ ID NO 4), and a cDNA sequence of Himalaya (SEQ ID NO 1). Predicted sequences were generated by identifying regions of the genomic sequences present in the Himalaya SSII cDNA. The ATG start codon and wild type stop codon are indicated, as are additional stop codons present in MK6827 (#) and 292 (&) respectively.

Figure 12 Comparison of amino acid sequences deduced from the genes encoding SSII from barley lines 292 (SEQ ID NO 7), Morex (SEQ ID NO 5), MK6827 (SEQ ID NO 8), Himalaya (SEQ ID NO 8). Additional stop codons in 292 and MK6827 are indicated by the symbols (&) and (#) respectively.

Figure 13. Position of the mutations in MK6827 (SEQ ID NO 2) and 292 (SEQ ID NO 4) in the barley SSII gene.

Figure 14. Development and use of a PCR assay for the 292 mutation. (a) schematic representation of an SSII region from Himalaya amplified by the primers ZLSS2P4 and ZLBSSIIP5 (b) representation of the region amplified from the SSII gene from 292 using ZLSS2P4 and ZLBSSIIP5, showing the absence of one NlaIV site (c) agarose gel electrophoresis of NlaIV digested products from barley; Lane M; DNA marker ladder, lane 1: MK6827, lane 2; Himalaya; lane 3, Tantangara; lane 4, 292; lane 5, 342.

Figure 15. SDS-PAGE electrophoresis of starch granule proteins. Panel (A) 8% Acrylamide (37.5:1 Acryl/Bis) SDS-PAGE gel, electroblotted and probed with a SSII antibody produced against purified granule-bound SSII protein from Wheat. (B) 12.5% acrylamide (30:0.135 Acryl/Bis), silver stained. The migration of molecular weight standards of defined mass (units are kd) are indicated on each side of the figure.

Figure 16. A schematic representation of DNA constructs designed to down regulate SSII expression following stable transformation of barley (1) The SSII gene from nucleotides 1 to 2972 (see Figure 9 for sequence) is inserted between the promoter and terminator in the sense orientation. (2) The SSII gene is inserted between the promoter and terminator in the anti-sense orientation from nucleotides 2972 to 1 (see Figure 9 for sequence). (3) Duplex construct in which intron 3 of the barley SSII gene (between nucleotides 1559 and 2851) of the Morex SSII genomic sequence is inserted between exons 2 and 3 from the barley SSII cDNA from Himalaya (nucleotides 363 to 1157 from Figure 9).

DETAILED DESCRIPTION OF THE INVENTION

Definitions

Glycaemic Index. Is a comparison of the effect of a test food such as white bread or glucose on excursions in blood glucose concentration. The Glycaemic Index is a measure of the likely

and resistant starch that does not, at least in broader forms of the invention require mixing of β glucan and soluble dietary fibre together or modification of the component parts.

To the best of the knowledge of the inventors the barley plant of the present invention is the first time that there has been a barley grain having elevated relative dietary fibre levels in the form of resistant starch having an elevated amylose level, that also has elevated levels of β glucan that are at the higher end of the typical levels of β glucan or that go beyond that level. Grains that have β glucan content that are still higher are of the waxy phenotype and therefore have low levels of amylose.

It is known that there is a wide variation in β glucan levels in barley in the range of about 4% to about 18% by weight of the barley, but more typically from 4% to about 8% (Izydorczyk *et al.*, (2000) *Journal of Agricultural and Food Chemistry* **48**, 982-989; Zheng *et al.*, (2000) *Cereal Chemistry* **77**, 140-144; Elfversson *et al.*, (1999) *Cereal Chemistry* **76**, 434-438; Andersson *et al.*, (1999) *Journal of the Science of Foods and Agriculture* **79**, 979-986; Oscarsson *et al.*, (1996) *J Cereal Science* **24**, 161-170; Fastnaught *et al.*, (1996) *Crop Science* **36**, 941-946). Enhanced barley strains have been developed, Prowashonupana for example, which have between about 15% and about 18% by weight β -glucan but has a waxy phenotype. This is sold commercially under the name Sustagrain™, (ConAgra™ Specially Grain Products Company, Omaha, Neb. USA).

The levels of β glucan contemplated by this invention may depend on the genetic background in which the amylopectin synthesis enzyme activity is reduced. However it is proposed that the reduction of the amylopectin synthesis activity will have the effect of elevating the relative level of dietary fibre which, in part, takes the form of amylose, and at the same time elevating the level of β glucan. One explanation for the concomitant elevation of β glucan with elevated relative amylose levels is that such elevation might be the result of a concentration effect of having reduced endosperm and may be further increased through the diversion of carbon from starch synthesis to β glucan synthesis.

Thus the grain of the barley plant preferably has a β glucan content that is greater than 6% of total non-hulled grain weight or more preferably greater than 7% and most preferably greater than 8%, however levels of β glucan in a waxy mutant has been measured as being as high as 15 to 18% and the present invention may contemplate levels as high, or higher, than that.

In a second preferable form the grain of the barley plant has a reduced gelatinisation temperature (as measured by differential scanning calorimetry) in addition to the relatively high amylose content. On the data shown for the exemplified barley this reduced gelatinisation temperature is not just reduced when compared to starch produced by barley with somewhat elevated amylose content but also when compared with starch produced from barley with starch having normal levels of amylose. Thus whilst the invention contemplates reduced gelatinisation temperatures relative to a corresponding high amylose starch, it may also contemplate a gelatinisation temperature reduced relative to that of starch with normal amylose levels.

Additionally in the genetic backgrounds thus far checked the starch is also characterised by a swelling in heated excess water that is lower than swelling of other starches tested.

In a third preferable form the starch has amylose levels of higher than 50% of the starch content which is a level never before found in unmodified starch derived from barley..

The starch of the present barley plant has a high relative amylose content and much higher than might be anticipated for a mutation in the SSII gene or other starch synthase gene. Thus in wheat mutants in SSII result in relative amylose levels of about 35% of starch. The amylose content of starch might be considered to be elevated when the content is significantly greater than the 25% or so that is present in normal barley grain and thus might be greater than about 30% w/w of total starch. Known barley plants considered to be high amylose have a content of 35-45%. The present invention however provides for barley with an amylose content that is greater than 50%, with is a level never before found in unmodified starch derived from barley.

The relative amylose content might be greater than 60% and more preferably, still greater than 70%. It may be desired to have even higher levels and thus it has been possible to achieve even higher levels in other plants by breeding with single mutations, such levels approach 90%. Thus the invention might encompass amylose levels of greater than 80% or greater than 90%.

It will be understood that the relative level of amylose referred to is in relation to total starch content, and thus the remainder of the starch might be predominantly of an intermediate type of starch or it might be predominantly amylopectin or a mixture of both. In the barley analysed the elevated level of amylose results from decreased amylopectin levels, and accordingly the relative level of amylose does not result from an increased synthesis of amylose.

It is known that β glucan has the effect of slowing digestion in the small intestine simply by its presence when together with another food component. Similarly it is known that resistant molecules that have close juxtaposition with starch granules help to mask the starch and contribute to its resistance by making it physically inaccessible. Elevated levels of amylose and other forms of starch as may arise from association with lipid will be further enhanced therefore by the presence and physical juxtaposition to the starch granules. Thus there is provided a significant enhancement of the effects of the resistant starch, as well as a provision of other beneficial effects arising from high β glucan levels.

Additionally it is known that there is a dose response in terms of the beneficial effects of resistant starch and β glucan. It is proposed therefore that the increased level of β glucan together with the increased levels of resistant starch will provide enhanced health benefits.

The combination of the levels of β glucan and resistant starch of at least preferred forms of this invention have not been found before and certainly not from one source without a degree of modification or purification and thus forms of the present invention provide for a single practical source of these benefits.

Another preferred aspect of the starch is that despite the high relative amylose content it also has a low gelatinisation temperature as measured by differential scanning calorimetry. This is in contrast with the general finding that high amylose starches tend to have a raised gelatinisation temperature which introduces restrictions on the manner in which high amylose starches can be utilised. On the data shown for the exemplified barley this reduced gelatinisation temperature is not just reduced when compared to starch produced by lines with somewhat elevated amylose content but also when compared with starch produced from barley with starch having normal levels of amylose. Thus whilst a preferred aspect of the invention contemplates reduced gelatinisation temperatures relative to corresponding high amylose starch it may also contemplate a gelatinisation temperature reduced relative to that of starch with

is about 3.75. Whereas the grains of the mutants and crosses examined are less than 3.2, preferably less than 3.0, but generally higher than about 2.

5 This low swelling gelatinisation characteristic is particularly useful where it is desired to increase the starch content of a food preparation, in particular a hydrated food preparation. In the present instance it might be desired to increase the dietary fibre content of a sol or other liquid preparation where there would otherwise be a restriction on delivery of the food preparation.

10 This characteristic in combination with the reduced gelatinisation temperature exhibited by the present starch provides a prospect of significantly enhancing the nutritional benefits of foods where there is a requirement of rapid preparation, such as instant soups and instant noodles.

15 It is postulated gelatinisation temperature effects are the result of an altered amylopectin structure in the endosperm of its grain, and one measurement of this structure is the distribution of chain lengths (degrees of polymerisation) of the starch molecules following debranching by isoamylase. An analysis of the chain length of the amylopectin content of the starch of the exemplified SSII mutants showed that when debranched they have a distribution of chain length in the range from 5 to 60 that is shorter than the distribution of starch yielded
20 by non-mutant lines upon debranching. Starch with shorter chain lengths will also have a commensurate increase in frequency of branching. Thus the starch may also have a distribution of shorter amylopectin chain lengths. The proportion of starch chains that have a degree of polymerisation that falls in the range of 6 to 11 residues may be greater than 25%, more preferably greater than 30% and most preferably greater than 35%. The proportion of
25 starch chains that have a degree of polymerisation that falls in the range of 12-30 residues may be less than 65%, more preferably less than 60% and most preferably less than about 55 %. The proportion of starch chains that have a degree of polymerisation that falls in the range of 31-60 residues may be less than about 10%, more preferably less than about 8% but also preferably greater than about 5% and more preferably greater than about 6%. Rather than
30 taken individually combination of proportions of the three chain length ranges might be taken as an indicator that a starch is of a type that accords with the present invention.

grain. This form of starch is usually associated with retrograded starch, in particular where there has been contact with lipids. In the case of the present invention it is postulated that the structure of the starch permits the formation of an intimate relationship between plant lipids and starch which results in the V-complex structure. It is thought that this form of starch may have health benefits because it has reduced digestibility and therefore may contribute to resistant starch.

Other forms of structure can also result from lipid-starch interaction and include non crystalline lipid-starch complexes. Thus the invention might also be said to reside in a barley plant exhibiting appreciable amounts of starch-lipid complexes in the starch content of the endosperm of its grain resulting from reduced levels of activity of one or more amylopectin synthesis enzymes. Starches that contain starch lipid complexes, including those that exhibit V-complex structure, are also usually resistant to digestion and thus contribute to the dietary fibre levels. Preferably the proportion of crystalline starch exhibiting a form of crystallinity characteristic of a starch-lipid complex is greater than about 50% and more preferably greater than about 80%.

The starch additional to the presence of the V-complex form of starch may also exhibit no appreciable amounts of A complex forms of starch. Absence of A-complex might be taken as indicator of the presence of a starch of this invention.

It is also found that the pasting temperature of strchs and product made from the grain of thisinvention are considerably elevated. The pasting temperatures in known starches is less than 70°C, and this is for both normal and high amylose starches. The starches of the present invention however preferably exhibit pasting temperatures of higher than about 75°C or more preferably higher than about 80°C. It will be noted that these are empirical measures and might be taken as relative to those measurement of the other starches.

The starch of the exemplified barley plant is found to have significant amounts of dietary fibre and resistant starch, presumably this increase is at least in part as a result of the high relative level of amylose, however there may also be a contribution of dietary fibre by reason of starch/lipid complexes, including V-complex, or because of the intimate associate of amylose

monocotyledonous plants such as barley and for regeneration of plants from protoplasts or immature plant embryos are well known in the art, see for example, Canadian Patent Application 2092588 by Nehra, Australian Patent Application No 61781/94 by National Research Council of Canada, Australian Patent No 667939 by Japan Tobacco Inc.,
5 International Patent Application PCT/US97/10621 by Monsanto Company, US Patent 5589617, and other methods are set out in Patent specification WO99/14314.

Other known approaches to altering the activity of the amylopectin synthesis enzyme, other than the use of mutations may also be adopted. Thus, for example, this could be by expression
10 of suitable antisense molecules that interfere with the transcription or processing of the gene or genes encoding the amylopectin synthesis enzyme. These might be based on the DNA sequence elucidated herein for the barley SSII gene. These antisense sequences can be for the structural genes or for sequences that effect control over the gene expression or splicing event. These sequences have been referred to above. Methods of devising antisense sequences are
15 well known in the art and examples of these are can be found in, for example, United States Patent 5190131, European patent specification 0467349 AI, European patent specification 0223399 AI and European patent specification 0240208, which are incorporated herein by reference to the extent that they provide methods for carrying out antisense techniques. Methods of introducing and maintaining such sequences in plants are also published and
20 known.

A variation of the antisense technique is to utilise ribozymes. Ribozymes are RNA molecules with enzymic function that can cleave other RNA molecules at specific sites defined by an antisense sequence. The cleavage of the RNA block the expression of the target gene.
25 Reference is made to European patent specification 0321201 and specification WO 97/45545.

Another molecular biological approach that might also be used is that of co-suppression. The mechanism of co-suppression is not well understood, but it involves putting an extra copy of a gene into a plant in the normal orientation. In some instances the additional copy of the gene
30 interferes with the expression of the target plant gene. Reference is made to Patent specification WO 97/20936 and European patent specification 0465572 for methods of implementing co-suppression approaches.

folate or antioxidants such as tocopherols and tocotrienols. Thus calcium is established in the provision of material for growth and deposition of bone and other calcified tissue and in lowering the risk of osteoporosis later in life. Folic acid is found to be protective against neural tube defects when consumed periconceptually and decreases the risk of cardiovascular disease thereby enhancing the effects of the combination of resistant starch and β -glucan. Folic acid also is thought to have an effect of lowering the risk of certain cancers. Tocopherol and tocotrienols carry the benefits of antioxidants and are believed to lower the risk of cancer and heart disease, and also have the effect of reducing the undesirable effects of oxidation of components of a food such as fatty acids which can result in rancidity. When these components of this preferred form of barley grain or products made therefrom constitute a convenient packaging with the one grain. One specific form of milled product might be one where the aleurone layer is included in the milled product. Particular milling process might be undertaken to enhance the amount of aleurone layer in the milled product. Such a method is referred to in Fenech *et al.*, ((1999) *J Nutr* 129:1114-1119). Thus any product derived from grain milled or otherwise processed to include aleurone layer and germ will have the additional nutritional benefits, without the requirement of adding these elements from separate sources.

It will be understood that the barley plant of the present invention is preferably one having grain that is useful for food production and in particular for commercial food production. Such a production might include making of flour or other product that might be an ingredient in commercial food production. A lower level of usefulness might be a starch content greater than about 12% or perhaps greater than about 15%. Or similarly this might include the capacity to mill the grain; thus whilst pearled barley may be produced from most forms of grain certain configurations of grain are particularly resistant to milling. Another characteristic that might have an impact on a variety producing a commercially useable grain is discolouration of the product produced. Thus where the husk or other portion of the grain exhibits significant colouration, for example purple, this will come through with the product and limits its commercial applications to niche applications such as being a component of a bread containing coloured whole or kibbled grains. It is generally also more convenient that the barley plants are naked, because the presence of husks on barley grains introduces greater difficulty in processing the grain. Another aspect that might make a barley plant of higher value is on the basis of starch extraction from the grain, the higher extraction rates being more useful. Grain shape is also another feature that can impact on the commercial usefulness of a

might be desired to make double mutations in other barley mutants available with shrunken endosperms where the causal gene is not known.

5 In a further aspect the invention could be said to reside in the grain produced from a barley plant as referred to in this specification.

It will also be understood that the invention encompasses a processed grain including a milled, ground, kibbled, pearled or rolled grain or product obtained from the processed or whole grain of the barley plant referred to above, including flour. These products may be then used in
10 various food products, for example farinaceous product such as breads, cakes biscuits and the like, or food additives, such as thickeners or to make malted or other barley drinks, noodles and quick soups.

Alternatively the invention encompasses starch isolated from the grain of the barley plant
15 referred to above. Starch might be isolated by known techniques.

It will be understood that one benefit of the present invention is that it provides for one or more products that are of particular nutritional benefit, and moreover it does so without the need to modify the starch or other constituents of the barley grain.
20

However it may be desired to make modifications to the starch, β glucan or other constituent of the grain, and the invention encompasses such a modified constituent.

The method of modification are those known, and include the extraction of the starch or
25 β glucan or other constituent by conventional methods and modification of the starches to for the desired resistant form.

Thus the starch or β glucan may be modified either singly or multiply though the use of a treatment selected from group including but not limited to, heat and/or moisture, physically
30 (for example ball milling), enzymatically (using for example α or β amylase, pullulanase or the like), chemical hydrolysis (wet or dry using liquid or gaseous reagents), oxidation, cross bonding with difunctional reagents (for example sodium trimetaphosphate, phosphorous oxychloride), or carboxymethylation.

While it is clear that at least these four activities are required for normal starch granule synthesis in higher plants, multiple isoforms of each of the four activities are found in the endosperm of higher plants and specific roles have been proposed for individual isoforms on the basis of mutational analysis (Wang *et al.*, 1998, Buleon *et al.*, 1998) or through the
5 modification of gene expression levels using transgenic approaches (Abel *et al.*, 1996, Jobling *et al.*, 1999, Schwall *et al.*, 2000). However, the precise contributions of each isoform of each activity to starch biosynthesis are still not known, and it is not known whether these contributions differ markedly between species. In the cereal endosperm, two isoforms of ADPglucose pyrophosphorylase are present, one form within the amyloplast, and one form in
10 the cytoplasm (Denyer *et al.*, 1996, Thorbjørnsen *et al.*, 1996). Each form is composed of two subunit types. The shrunken (*sh2*) and brittle (*bt2*) mutants in maize represent lesions in large and small subunits respectively (Girouiz and Hannah, 1994). Four classes of starch synthase are found in the cereal endosperm, an isoform exclusively localised within the starch granule, granule-bound starch synthase (GBSS), two forms that are partitioned between the
15 granule and the soluble fraction (SSI, Li *et al.*, 1999a, SSII, Li *et al.*, 1999b) and a fourth form that is entirely located in the soluble fraction, SSIII (Cao *et al.*, 2000, Li *et al.*, 1999b, Li *et al.*, 2000). GBSS has been shown to be essential for amylose synthesis (Shure *et al.*, 1983), and mutations in SSII and SSIII have been shown to alter amylopectin structure (Gao *et al.*, 1998, Craig *et al.*, 1998). No mutations defining a role for SSI activity have been
20 described.

Three forms of branching enzyme are expressed in the cereal endosperm, branching enzyme I (BEI), branching enzyme IIa (BEIIa) and branching enzyme IIb (BEIIb) (Hedman and Boyer, 1982, Boyer and Preiss, 1978, Mizuno *et al.*, 1992, Sun *et al.*, 1997). In maize and rice, high
25 amylose phenotypes have been shown to result from lesions in the BEIIb gene (Boyer and Preiss, 1981, Mizuno *et al.*, 1993). In these mutants, amylose content is significantly elevated, and the branch frequency of the residual amylopectin is reduced. In addition, there is a significant pool of material that is defined as "intermediate" between amylose and amylopectin (Boyer *et al.*, 1980, Takeda, *et al.*, 1993). Mutations defining the roles of BEIIa
30 and BEI have yet to be described, although in potato down regulation of BEI alone causes minimal affects on starch structure (Filipse *et al.*, 1996). However, in potato the combination of down regulation of BEII and BEI provides a much higher amylose content than the down-regulation of BEII alone (Schwall *et al.*, 2000). Two types of debranching enzymes are present in higher plants and are defined on the basis of their substrate specificities, isoamylase

starch biosynthesis and illustrate how mutations in specific genes can have differing impacts on starch structure from one species to another.

Materials and Methods

5 *Mutagenesis and Screening*

The hull-less barley variety "Himalaya" was mutagenised using sodium azide according to Zwar and Chandler (1995). Selection of variants with altered grain morphology was carried out according to Green *et al.*, (1997). A total of 75 lines with shrunken endosperm phenotypes were identified and maintained according to Green *et al.*, (1997).

10

Starch Isolation

Starch was isolated from barley grain using the method of Schulman *et al.* (1991).

Methods for Amylose Determination

- 15 Determinations of the amylose/amylopectin ratio by an HPLC method for separating debranched starches, and an iodine binding method, were carried out as described by Batey and Curtin, (1996). Analysis of the amylose/amylopectin ratio by the analysis on non-debranched starches was carried out according to Case *et al.*, (1998).

20 *Starch Content Measurement*

Starch was determined using the total starch analysis kit supplied by Megazyme (Bray, Co Wicklow, Republic of Ireland).

Protein Content

- 25 Nitrogen was determined by the Kjeldahl method, and protein contents were calculated using a factor of 5.7.

β -Glucan Levels

- β -Glucan was determined using the kit supplied by Megazyme (Bray, Co
30 Wicklow, Republic of Ireland).

Starch Chain Length Distribution

Doubled haploids were produced from F1 plants derived from crosses between 292 and *Hordeum vulgare* cv Tantangara, and between 342 and *H. vulgare* cv Tantangara by Dr P. Davies, Waite Institute, Adelaide, Australia.

5 *Linkage Analysis*

Genetic linkage data was calculated using MapManager.

Construction of barley cDNA library

Five mgs of polyA+ mRNA from 10, 12 and 15 days post-anthesis of barley endosperm tissues was used for cDNA synthesis according to the protocols (Life Technology). The *NotI*-(dT)18 primer (Pharmacia Biotech) was used for the first stand of cDNA synthesis. The double strand cDNAs were ligated with a *SalI*-*XhoI* adapter (Stratagene) and cloned to the *SalI*-*NotI* arms of ZipLox (Life Technology) after digestion of cDNAs with *NotI* followed by size fractionation (SizeSep 400 spun Column of Pharmacia Biotech). The ligated cDNAs were packaged with Gigapack III Gold packaging extract (Stratagene). Titre of the library was 2×10^6 pfu tested with Y1090(ZL) strain of *E.coli*.

Cloning of specific cDNA regions of barley starch synthase II using PCR

The cDNA clone, wSSIIp1, was used for the screening of a cDNA library of barley. The cDNA clone, wSSIIp1 was generated by PCR using the primers ssIIa (TGTTGAGGTTCC ATGGCACGTTC SEQ.ID. NO 9) and ssIIb (AGTCGTTCTGCCGTATGATGTCG SEQ. ID. NO 10), amplifying the region between nucleotide positions 1,435 and 1,835 of wSSIIA (GenBank accession no: AF155217).

The amplification was performed using a FTS-1 thermal sequencer (Corbett, Australia) for 1 cycle of 95°C for 2 minutes; 35 cycles of 95°C for 30 seconds, 60°C for 1 minutes, 72°C for 2 minutes and 1 cycle of 25°C for 1 minute. The fragment wSSIIp1 was cloned into a pGEM-T vector (Promega)

Screening of barley cDNA library

A cDNA library, constructed from RNA from the endosperm of barley cv Himalaya, was screened with a 347-bp cDNA fragment, wSSIIp1 at the hybridisation conditions as previously described (Rahman *et al.*, 1998). Hybridisation was carried out in 50% formamide, 6 x SSPE, 0.5% SDS, 5 x Denhardt's and 1.7 µg/mL salmon sperm DNA at 42°C for 16 h, then washed 3 x with 2 x SSC containing 0.1% SDS at 65°C for 1 h per wash.

Doubled haploids were produced from F1 plants derived from crosses between 292 and *Hordeum vulgare* cv Tantangara, and between 342 and *H. vulgare* cv Tantangara by Dr P. Davies, Waite Institute, Adelaide, Australia.

5 *Backcrossing Strategy*

Crosses were made between 292 and *Hordeum vulgare* cv Sloop to generate F1 seed. Plants derived from the F1 seed were selfed to generate a population of F2 seed. The plants growing from these F2 seed were tested using a PCR assay and plants homozygous for the 292 mutation were backcrossed to Sloop (BC1). The F1 plants resulting from BC1 were again
10 tested by PCR and plants heterozygous for the 292 mutation selected, and crossed back to Sloop (BC2). The F1 plants derived from BC2 were again analysed by PCR and plants heterozygous for the 292 mutation selected. These plants were either selfed to generate a BC2F2 population, or crossed again to Sloop (BC3). The F1 plants derived from BC3 were again analysed by PCR and plants heterozygous for the 292 mutation selected. These plants
15 were selfed to generate a BC3F2 population. Plants derived from these seed were tested by PCR and plants homozygous for the 292 mutation selected for single seed descent and seed increase.

Results

20 *Selection of Mutants*

The identification of a range of mutants in the hull-less or naked barley variety "Himalaya" induced by a sodium azide treatment has been previously reported by Zwar and Chandler (1995). A group of 75 shrunken grain mutants were identified by the inventors and the amylose content of the starch from the shrunken seed was determined by HPLC (Figure 1).
25 Two lines, 292 and 342, were found to have amylose contents of 71 and 62.5% respectively (Table 1). The amylose contents of 292 and 342 were substantially higher than the previously well characterised AC38 line (47% amylose, see Table 1). This study defines the genetic basis of the novel high amylose phenotype displayed by 292 and 342, and describes effects of the causal mutation on grain and starch structure and functionality.

30

Grain Characteristics

Grain size and morphology:

The effects of the mutation on grain weight and morphology are marked (Table 2). The grain weight is reduced from 51 mg for the parent line Himalaya, to 32 mg for 292 and 35 mg for

mg/caryopsis in 292 and 4.8 mg/caryopsis in 342. In contrast, there is a dramatic reduction in amylopectin synthesis per caryopsis, from 18.7 mg in Himalaya, to 1.6 mg in 292 and 2.9 mg in 342.

5 *Chain Length Distribution*

The chain length distribution of the starch following isoamylase debranching was carried out using fluorophore-assisted carbohydrate electrophoresis (FACE). The chain length distribution of the 292 and 342 mutants, and Himalaya, are shown in Figure 3a. Figure 3b shows a difference plot in which the normalised chain length distributions for the 292 and 342 mutants are subtracted from the normalised distribution of Himalaya. The percentages of chain lengths from DP 6-11, DP 12-30 and DP 31-65 have been calculated and are presented in Table 3. There is a marked shift in the 292 and 342 mutants in chain length distribution such that there is a higher percentage of chains in the region from DP6-11 compared to DP12-30.

15

Differential Scanning Calorimetry

The gelatinisation temperature of the mutants was investigated using differential scanning calorimetry, and the data is shown in Table 4. Both 292 and 342 yield starches that have markedly lower gelatinisation temperatures than the Himalaya starches, with respect to onset, peak and final temperatures for the gelatinisation peak. The enthalpy for the gelatinisation peak for the 292 and 342 mutants is also dramatically reduced in comparison to the wild type. The amylose/lipid peak onset temperature is also reduced for the 292 and 342 mutants, however, the enthalpy is increased, consistent with the increased amylose content of the mutants.

25 *Starch Viscosity by RVA*

RVA analysis of barley wholemeal samples was conducted in order to examine their pasting viscosity. Previous studies have shown that analysis of wholemeal samples is strongly correlated with the analysis of isolated starches (Batey *et al.*, 1997). The analysis showed that there are major differences between the barley genotypes studied (see Table 5 and Figure 4). Two barley varieties containing wild type starch, Himalaya and Namoi, showed typical RVA profiles in which there was a prominent peak viscosity, followed by a decline in viscosity to a holding strength, followed by an increase in viscosity as the temperature is reduced to a final viscosity. As is generally observed for barley starches, the final viscosities for the wild type starches were equivalent to, or less than, the peak viscosities (Table 5). In AC38, a prominent peak viscosity was obtained, however, because of the elevated amylose content of this line, the

Dietary Fibre

Dietary fibre analysis was conducted according to the AOAC procedure and showed that there was an increase in dietary fibre in 292 and 342, and that this increase in dietary fibre was due to an increase in insoluble dietary fibre rather than soluble dietary fibre (Table 1), consistent with components of the dietary fibre being resistant starch and β -glucan. It is to be noted that this measure of dietary fibre is a chemically determined one which is quite distinct from the physiological measure relevant from a nutritional point of view.

10 Genetic Basis of the Mutation*Segregation ratio*

Crossing of the mutation to barley varieties not displaying the shrunken endosperm phenotype of 292 or 342 demonstrated that the mutation is a straightforward recessive mutation, displaying a 3 normal : 1 shrunken ratio in the F2 seed of outcrossed populations, and 1 normal : 1 shrunken ratio in the seed of a doubled haploid population developed following a single outcross (see Table 6). Normal is defined as seed with an L/T ratio of <3.5 , shrunken seed as seed with an L/T ratio of >3.5 .

Allelic nature of mutants

20 The 292 and 342 mutations were shown to be allelic through the analysis of progeny from crosses of 292 and 342. All F1 seed derived from reciprocal crosses showed grain weight and grain morphology phenotypes within the range of sizes and shapes observed for the parental 292 and 342 lines, and outside of the range of seed size and shape found for the parental Himalaya line. Furthermore, all F2 seed derived from 292 x 342 F1 plants showed the typical shrunken seed phenotype of the 292 and 342 mutants.

Analysis of the grain morphology and starch characteristics of a series of shrunken grain mutants available from the Barley Germplasm Collection (USDA-ARS, National Small Grains Germplasm Research Facility, Aberdeen, Idaho 83210, USA) suggested that the line MK6827 (BGS31, also referred to as GSHO 2476), carrying the *sex6* mutation showed a highly similar set of starch and grain characteristics to the 292 and 342 mutations. Crosses were established between 292 and MK6827 and all F1 grain showed the typical 292 phenotype with respect to grain weight and shrunken seed phenotype. F2 seed derived from the 292 x MK6827 F1 plants all showed shrunken endosperm phenotype with L/T ratios of >4 . In contrast, F2 seed

mapped within 16.3 cM of the *nud* locus. This location is consistent with previous mapping data for the allelic *sex6* mutation (Netsvetaev, 1990, Netsvetaev and Krestinkov, 1993, Biyashev *et al.*, 1986, Netsvetaev, 1992).

5 *Identification of the causal gene*

The *nud* gene has been demonstrated to be located on barley chromosome 7H (Figure 8, Fedak *et al.*, 1972). In wheat, three starch synthases (GBSS, SSI and SSII), and an isoamylase-type debranching enzyme (S. Rahman, personal communication) are located on the short arm of chromosome 7, the homologous chromosome (Yamamori and Endo, 1996, Li *et al.*, 1999a, Li *et al.*, 1999b, Li *et al.*, 2000). The close linkage to the *nud* locus suggested that the most probable candidate gene was the SSII gene. The wheat SSII gene has been cloned at the cDNA level (Li *et al.*, 1999b; Genbank Accession No. AF155217) and at the genomic level (Li *et al.*, personal communication), and a barley cDNA has been isolated and cloned (Figure 9). The sequencing of barley and wheat SSII genomic sequences shows that the genes have very similar exon/intron structures, however, the lengths of the intron regions differ between sequences (Figure 10). Comparison of the Morex genomic sequence and the sequence of a cDNA from Himalaya (Figure 9) lead to the identification of deduced cDNA sequences from Morex, 292 and MK2827.

20 A G to A transition mutant was found in the SSII gene from 292 at a position that corresponds to 1829 of the alignment shown in Figure 11. This mutation introduces a stop codon into the 292 SSII open reading frame (Figure 12). Sequence analysis of Tantangara and Himalaya showed that both wild type genes were identical in this region and both 292 and 342 contained the same G to A transition mutation. The introduced stop codon would truncate the gene product such that the entire C-terminal catalytic domain of the starch synthase II gene would not be translated, and it is therefore highly likely that all SSII activity is abolished by this mutation.

30 A G to A transition was also present in MK6827, at position 242 of the alignment shown in Figure 11 and the Himalaya cDNA sequence in Figure 9. This mutation also introduces a stop codon into the 292 SSII open reading frame (Figure 12) and would prevent translation of over 90% of the SSII gene, abolishing SSII activity encoded by this gene.

DNA marker status is given in Table 8. More comprehensive analysis of the composition of these lines is given in Table 9, including RVA analysis, β -glucan content and flour swelling volume. The data shows that the lines carrying the 292 mutation have significantly different RVA parameters (as exemplified by the Peak/Final Viscosity ratio), higher β -glucan content, and altered flour swelling volumes.

In the second example, the mutation was transferred by performing two backcrosses from 292 to a cultivar with normal starch properties (cv Sloop). The F2 seed from three backcross 2 F1 plants was collected for analysis. The F2 seed were categorized into seed with an L:T ratio of >3.5 and an L:T ratio of <3.5 . The distribution of seeds between these classes was consistent with expectations for a single recessive gene. Flour swelling volume data for the categories of seeds derived from each plant are shown in Figure 10 and shows that the starch swelling trait was clearly transferred through the breeding process into lines with an average of 75% Sloop background.

Discussion

We describe the isolation of novel mutants, 292 and 342, in barley that have a shrunken endosperm phenotype. Analysis of grain composition demonstrates that the shrunken phenotype is due to a significant decrease in starch content, and analysis of starch composition shows that this decrease is manifested as a high amylose phenotype that arises because of a decrease in amylopectin synthesis.

The 292 and 342 mutants possess a unique combination of grain and starch properties, in containing both increased β -glucan levels and resistant starch. The β -glucan levels of the lines are increased approximately 15% above that expected by the effect of reduced starch content, suggesting that carbon unable to be converted to starch is diverted to β -glucan synthesis. Determinations of dietary fibre levels demonstrate that the grain from the mutants have increased levels of dietary fibre, and that this increase is due to an increase in insoluble dietary fibre.

This combination of properties indicates that these mutants may have very interesting potential as components of the human diet. First, the elevated β -glucan levels suggests that the lines may be useful in lowering cholesterol through the well established action of β -glucan in reducing cholesterol levels. Secondly, the presence of resistant starch indicates that the lines

the crystal form shifts from the A type typical of cereal starches to a mixture of V and B types. The V type is typical of amylose and reflects the amylose component of the starch complexed with fatty acids, while the B form is derived from amylopectin and presumably reflects the residual amylopectin content of the starch (Buleon *et al*, 1998).

- 5
- Analysis of the genetic basis of the 292 and 342 mutations demonstrates that the mutations are simple recessive mutations that give typical Mendelian ratios in outcrossing experiments. Crossing studies demonstrated that 292 and 342 are allelic. Further analysis of the interaction between 292 and other shrunken endosperm mutations in crossing experiments demonstrated
- 10 that the 292/342 mutations were also allelic with the Sex6 mutation in the line MK6827. This mutation has previously been mapped and shown to be located within 3 cM of the centromere on the short arm of chromosome 7H (Netsvetaev, 1990, Netsvetaev and Krestinkov, 1993, Biyashev *et al.*, 1986, Netsvetaev, 1992).
- 15 A doubled haploid population between the husked barley Tantangara and the naked 292 mutant was established and the shrunken endosperm mutation mapped to the short arm of chromosome 7HS, to within 16 cM of the *nud* gene, a location consistent with the map location of the Sex6 mutation.
- 20 The localisation of the gene to the region adjacent to the centromere on the short arm of chromosome 7HS demonstrates that the causal mutation (*sex6*) is in a different gene to the mutation that causes the high amylose phenotype in AC38 (*amol1*) which has been mapped to chromosome 1H (Schondelmeier *et al* 1992). The map location suggested that one candidate for the gene disrupted in the *sex6*/292 mutation was starch synthase II, known in wheat to be
- 25 localised in the same region of the chromosome (Yamamori and Endo 1996, Li *et al*, 1999b). Sequence analysis of the 292 and 342 mutants showed that there was a G to A transition mutation in the gene which would cause truncation of the gene such that the C-terminal region containing the active site of the enzyme would not be translated, presumably leading to the synthesis of a completely inactive protein. Furthermore, the sequencing of the SSII gene from
- 30 MK6827 showed a G to A transition mutation at position 242 which would also cause truncation of the gene. This result confirms the allelic nature of the 292 and MK6827 mutations.

The availability of the sequence of the SSII gene and barley transformation systems provides the tools required to knock out the SSII gene using gene suppression technologies, in order to produce a comparable phenotype to that found with the SSII mutations. A recently developed highly effective strategy is to produce a hairpin construct designed to produce a double stranded RNA which would suppress the endogenous SSII activity. While complete knock out mutants analogous to the mutations described here would be of interest, the use of DNA constructs with differing promoters, and the recovery of transgenes with differing levels of hairpin construct expression, would allow the impact of titrating the expression of the gene from normal levels to complete knockdown levels to be assessed.

The mutations were shown to be able to be transferred from 292 into alternative barley genetic backgrounds, while retaining essential features of the original 292 mutation. In Tables 9 and 10, phenotypic data for 292 x Tantaranga doubled haploid progeny, and the seed from a second backcross to Sloop, are shown, and indicate that the phenotypes are transferred through the breeding process.

Table 1
Barley Grain Composition

	Starch Content (%) ^a	Amylose Content By HPLC (%) ^b	Amylose Content by iodine binding (%)	Protein Content (%) ^a	β-glucan (%) ^a	Total Dietary Fibre ^a (%)	Insoluble Dietary Fibre ^a (%)	Soluble Dietary Fibre ^a (%)
Glacier	n.d.	31.0	n.d.	11.5	4.3	21.6	16.6	5
AC38	47	47.4	60.6	10.4	5.8	24.9	28.8	6.1
Himalaya	49	25	25.4	10.0	4.8	27.1	18.1	9
292	17.7	71	68.9	15.0	9.5	30.3	21.4	8.9
342	21.9	62.5	71.7	15.7	8.3	28.3	19.4	8.9
MK6827	10.2	n.d.	44.4	21.3	n.d.	n.d.	n.d.	n.d.
Waxiro	42.8	n.d.	5.0	14.6	n.d.	19.8	12.7	7.1
Tantaranga	51.6	n.d.	29.5	14.6	n.d.	17.2	12.7	4.5

^a % grain weight, 14% moisture

^b % of total starch content

n.d. not determined

Table 4
Barley Starch Thermal Properties Measured by DSC

	Peak 1				Peak 2			
	Onset	Peak	End	ΔH	Onset	Peak	End	ΔH
Glacier	55.4	59.3	65.3	4.2	93.9	101.4	107.7	0.87
AC38	55.0	62.2	68.2	3.9	89.3	100.1	106.9	1.195
Himalaya	56.8	60.9	68.0	4.5	93.1	101.8	108.3	0.78
292	46.0	51.2	58.1	0.29	88.7	97.7	104.9	1.34
342	45.2	50.4	56.8	0.47	86.5	97.0	105.0	1.59

5

Table 5
RVA Parameters for Barley Starches

	Peak Viscosity	Breakdown	Holding Strength	Setback	Final Viscosity	Normalised Final Viscosity*	Pasting Temp (C)
Himalaya	871.5	653.1	218.4	235.8	454.2	926	64.9
Namoi	621.7	367.5	254.2	375.3	629.5	1284	65.9
AC38	226.7	87.3	139.4	188.4	327.8	697	68.9
292	92.1**	***	133.9	230	363.9	2055	89.5
342	110.9**	***	144.9	264.5	409.4	1869	87.9
MK6827	18.2**	***	25.7	43.3	69	676	n.d.

* Final viscosity divided by starch content of wholemeal
 ** Value registered at time of peak viscosity for Himalaya
 *** Value was less than zero
 n.d. not determined

10

15

Table 6
Starch Crystallinity Data

Sample	% H ₂ O (W.B)	Crystallinity %*	A %*	B %*	V %*
292	29.6	9	-	13	87
342	35.8	12	-	18	81
AC38	26.1	19	93	7	(traces)
Himalaya	27.7	27	93	7	(traces)
Waxiro	29.7	41	94	6	-

(* $\pm 5\%$)

20

Table 7
Progeny Analysis

Table 8
Scoring of 292 x Tantangara Doubled Haploid Lines

Line Number ^a	Husk ^b	Seed Weight (mg)	L/T Ratio ^c	DP6-11 (%) ^d	Amylose Content ^e	PCR ^f
1	N	26	3.8	35.87	50.2	292
2	N	24	4.21	36.87	56.2	292
3	H	43	3.32	25.45	18.3	Wt
5	N	40	4.58	39.47	55.5	292
7	N	34	4.28	19.63	43.0	292
8	H	48	3.02	21.6	46.7	Wt
9	N	31	2.76	22.89	25.9	Wt
10	N	26	3.02	27.56	21.1	Wt
11	N	34	3.55	37.90	44.7	292
12	H	50	2.94	26.37	32.8	Wt
13	N	27	4.29	38.68	48.4	292
14	H	56	3.07	22.98	20.8	Wt
15	H	46	2.74	24.88	22.9	Wt
16	H	43	2.78	25.40	18.3	Wt
17	N	31	3.8	37.37	54.2	292
18	N	31	4.51	37.46	57.5	292
19	H	26	3.1	29.57	22.7	Wt
20	H	53	3.04	25.42	23.8	Wt
21	N	31	4.5	38.51	59.1	292
22	N	27	4.63	37.25	27.2	292
23	H	47	2.73	24.11	21.2	Wt
24	N	27	4.58	36.89	42.0	292
26	H	35	3.57	19.50	15.1	Wt
27	H	22	4.3	36.81	48.6	292
28	N	31	4.34	38.88	37.0	292
30	N	30	4.04	38.05	48.4	292
31	N	23	4.25	37.07	51.7	292
32	H	48	2.62	20.67	13.0	Wt
33	N	25	4.92	35.68	33.3	292
34	N	31	4.01	38.34	46.1	292
35	H	43	3.16	20.07	23.6	Wt
36	N	26	4.33	36.93	29.7	292
38	H	38	3.01	21.11	9.1	Wt
39	H	33	2.92	20.49	23.5	Wt
40	H	36	2.99	19.57	2.2	Wt
41	N	30	4.05	37.82	40.9	292
42	H	47	2.95	20.80	11.9	Wt
43	N	40	3.24	21.97	18.1	Wt
45	H	52	2.78	19.97	14.5	Wt
46	N	29	4.44	35.87	32.1	292
47	N	35	3.69	36.34	92.9	292
48	H	31	2.54	20.27	13.4	Wt
49	H	54	2.94	22.29	19.3	Wt
50	H	50	2.94	21.92	20.6	Wt

^c L/T ratio: length to thickness ratio

^d Percentage of chains in debranched starch with DP6 to DP11, calculated on a molar basis as a percentage of chains eluting between DP6 and DP65

^e Amylose content determined by iodine blue value

- 5 ^f PCR score. 292, PCR reaction yields band which yields 169 bp band plus 103 bp on NlaIV digestion; Wt, PCR reaction yields band which yields 111 bp, 103 bp and 57 bp band on NlaIV digestion

Table 10
Flour Swelling Data for BC2F2 seed

Line	Swelling Volume
C5/1 Plant 1 L:T>3.5	2.118
C5/1 Plant 1 L:T<3.5	6.913
65/2 Plant 1 L:T>3.5	2.382
65/2 Plant 1 L:T<3.5	7.565
65/2 Plant 2 L:T>3.5	2.409
65/2 Plant 2 L:T<3.5	6.707

- Akerberg *et al.*, (1998). *Journal of Cereal Science* **28**, 71-80.
- Andersson *et al.*, (2000). *Cereal Chemistry* **77**, 463-467
- Andersson *et al.*, (1999). *Journal of the science of food and agriculture*. **79**, 979-986.
- Andreev *et al.*, (1999). *StarchStarke*. **51**, 422-429.
- 5 Banks *et al.*, (1971). *Starch*. **23**, 12-15.
- Batey and Curtin. (1997). *Starch* **48**, 338-344.
- Batey *et al.*, (1997). *Cereal Chemistry* **74**, 497-501.
- Bergh *et al.*, (1999). *Animal Feed Science and Technology* **78**, 215-226.
- Bhatty (1999). *Cereal Chemistry* **76**, 589-599.
- 10 Biyashev *et al.*, (1986). *Soviet Genetics* **22**, 296-303.
- Boyer and Preiss, (1978). *Carbohydrate Research* **61**, 321-334.
- Boyer and Preiss, (1981). *Plant Physiology* **67**, 1141-1145.
- Boye *et al.*, (1980). *Starch* **32**, 217-222.
- Buleon *et al.*, (1998). *International Journal of Biological Macromolecules* **23**, 85-112.
- 15 Calvert *et al.*, (1976). *Nutrition Reports. International* **14**, 55-61.
- Campbell *et al.*, (1994). *Cereal Chemistry* **71**, 464-468.
- Cao *et al.*, (2000). *Archives. of Biochemistry and Biophysics*. **373**, 135-146.
- Case *et al.*, (1998). *Journal of Cereal Science* **27**, 301-314.
- Castillo, *et al.*, (1994). *Bio/technology* **12**, 1366-1371.
- 20 Cho, *et al.* (1999). *Plant Science (Limerick)* **148**, 9-17.
- Craig *et al.*, (1998). *Plant Cell* **10**, 413-426.
- Czuchajowska *et al.*, (1998). *Cereal Chemistry* **75**, 747-754.
- Czuchajowska *et al.*, (1992). *Cereal Chemistry* **69**, 413-418.
- Denyer *et al.*, (1996). *Plant Physiology* **112**, 779-785.
- 25 Elfverson *et al.*, (1999). *Cereal Chemistry* **76**, 434-438.
- Eslick and Ries, (1976). *Barley Genetics Newsletter* **6**, 21-22.
- Fastnaught, *et al.*, (1996). *Crop Science* **36**, 941-946.
- Fedak *et al.*, (1972). *Canadian Journal of Genetics and Cytology* **14**, 949-957.
- Filpse *et al.*, (1996). *Planta* **198**, 340.
- 30 Fontaine *et al.*, (1993). *Journal of Biological Chemistry* **268**, 16223-16230.
- Fujita *et al.*, (1999) *Breeding. Science* **49**, 217-219.
- Fuwa *et al.*, (1999). *Starch/Starke*. **51**, 147-151.
- Gao *et al.*, (1998). *Plant Cell* **10**, 399-412.
- Giroux and Hannah. (1994). *Molecular and General Genetics* **243**, 400-408.
- 35 Gless, *et al.* (1998). *Journal of Plant Physiology* **152**, 151-157.

- Persson *et al.*, (1996). *Sveriges. Utsadesforenings. Tidskrift* **106**, 79-86.
- Pomeranz, (1992). *European. Journal of Clinical. Nutrition* **46**, S63-S68.
- Pomeranz *et al.*, (1972). *Cereal Chemistry* **49**, 629-635.
- Rahman *et al.*, (1995). *Australian Journal of Plant Physiology* **22**, 793-803.
- 5 Ramage and Eslick, (1975). *Barley Genetics Newsletter* **5**, 114.
- Ramage, and Eslick, (1975). *Barley Genetics Newsletter* **6**, 115.
- Schondelmaier *et al.*, (1992). *Plant Breeding* **109**, 274-280.
- Schulman and Kammiovirta, (1991). *Starch* **43**, 387-389.
- Schwall *et al.*, (2000). *Nature Biotechnology* **18**, 551-554.
- 10 Shure *et al.*, (1983). *Cell* **35**, 225-233.
- Somers *et al.*, (1992). Fertile, transgenic oat plants. *Bio/technology* **10**, 1589-1594.
- Somers, *et al.* (1994). Genetic engineering of oat. 37-46.
- Song and Jane, (2000). *Carbohydrate polymers* **41**, 365-377.
- Sun *et al.*, (1997). *The New Phytologist* **137**, 215-215.
- 15 Sundberg *et al.*, (1998). *Journal of the Science of Food and Agriculture*. **76**, 457-463.
- Swanston, (1992). *Barley Genetics Newsletter* **22**, 66-67.
- Swanston *et al.*, (1995). *Journal of Cereal Science* **22**, 265-273.
- Takeda *et al.*, (1993). *Carbohydrate Research* **246**, 273-281.
- Thorbjornsen *et al.*, (1996). *Plant Journal* **10**, 243-250.
- 20 Tingay, *et al.*, (1997). *Plant Journal* **11**, 1369-1376.
- Topping, (1999). *Asia Pacific Journal of Clinical Nutrition* **8**, S22-S26.
- Topping *et al.*, (1997). *The Journal of Nutrition* **127**, 615-615.
- Vasanthan and Bhatt, (1995). *Cereal Chemistry* **72**, 379-384.
- Vasanthan and Bhatt, (1998). *Starch-Starke*. **50**, 286-291.
- 25 Walker and Merritt, (1968). *Nature* **221**, 482-484.
- Wan and Lemaux, (1994). *Plant Physiology* **104**, 37-48.
- Wang *et al.*, (1998). *Journal of Experimental Botany* **49**, 481-502.
- Wang, *et al.*, (1993). *Cereal Chemistry* **70**, 521-525.
- Wilson and McNab, (1975). *British Poultry Science* **16**, 497-504.
- 30 Xue *et al.*, (1996). *Cereal Chemistry* **73**, 588-592.
- Yamamori, (1998). (Anon.), pp. 300-302. University Extension Press, University of Saskatchewan, Saskatoon.
- Yamamori and Endo, (1996). *Theoretical and Applied Genetics* **93**, 275-281.
- Yoshimoto *et al.*, (2000). *Cereal Chemistry* **77**, 279-285.
- 35 Zhang, *et al.* (1999). *Plant Cell Reports*. **18**, 959-966.

CLAIMS

1. Starch obtained from starch granules of grain of a barley plant the barley plant having a reduced level of SSII activity, said starch granules having a high amylose content
5 by reason of a reduced amylopectin content.
2. The starch of claim 1 with also exhibiting low gelatinisation temperatures.
3. The starch of claim 2 wherein the onset of the first peak detected by differential
10 scanning calorimetry is reduced.
4. The starch of claim 2 wherein the first peak detected by differential scanning calorimetry is reduced.
- 15 5. The starch of claim 2 wherein the enthalpy (ΔH) of the first peak is reduced.
6. The starch of claim 2 exhibits a low swelling volume.
7. The starch of claim 6 having a swelling volume of less than about 3.2.
20
8. The starch of claim 6 having a swelling volume of less than about 3.0.
9. The starch of claim 6 having a swelling volume of higher than about 2.0.
- 25 10. The starch of claim 1 wherein the starch when gelatinised exhibits reduced peak viscosity.
11. The starch of claim 1 wherein the pasting temperature of the starch is higher than
30 80°C
12. The starch of claim 1 wherein the pasting temperature of the starch is higher than
75°C
- 35 13. The starch of claim 1 wherein amylose levels in the grain are higher than 30% (w/w) of the starch content.

25. The starch of claim 23 wherein the proportion of starch that exhibits crystallinity is less than about 20% (
- 5 26. The starch of claim 1 wherein the starch exhibits reduced amylopectin chain length distribution
27. The starch of claim 26 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 6 to 11 residues is greater than 25%
- 10 28. The starch of claim 26 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 6 to 11 residues is greater than 30%
29. The starch of claim 26 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 6 to 11 residues is greater than 35%
- 15 30. The starch of claim 26 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 12-30 residues is less than 65%,
31. The starch of claim 26 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 12-30 residues is less than 60%,
- 20 32. The starch of claim 26 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 12-30 residues is less than 55%,
- 25 33. The starch of claim 26 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 31-60 residues is less than about 10%.
34. The starch of claim 26 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 31-60 residues is less than about 8%.
- 30 35. The starch of claim 34 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 31-60 residues is greater than about 5%.
- 35 36. The starch of claim 34 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 31-60 residues is greater than about 6%.

49. The grain of claim 48 wherein the pasting temperature of the starch is higher than 75°C
- 5 50. The grain of claim 49 wherein the pasting temperature of the starch is higher than 80°C
51. The grain of claim 37 wherein the grain has an elevated level of β glucan.
- 10 52. The grain of claim 51 wherein the β glucan content that is greater than 6% of total non-hulled grain weight
53. The grain of claim 51 wherein the β glucan content that is greater than 7% of total non-hulled grain weight
- 15 54. The grain of claim 51 wherein the β glucan content that is greater than 8% of total non-hulled grain weight
55. The grain of claim 51 wherein the β glucan content that is greater than about 15% of total non-hulled grain weight
- 20 56. The grain of claim 37 wherein the amylose content is higher than 30% (w/w) of the starch content.
57. The grain of claim 37 wherein the amylose content is higher than 50% (w/w) of the starch content
- 25 58. The grain of claim 37 wherein the amylose content is higher than 60% (w/w) of the starch content
- 30 59. The grain of claim 37 wherein the amylose content is higher than 70% (w/w) of the starch content
60. The grain of claim 37 exhibit appreciable amounts of starch associated lipid

72. The grain of claim 69 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 6 to 11 residues is greater than 35%.

5 73. The grain of claim 69 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 12-30 residues is less than 65%.

74. The grain of claim 69 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 12-30 residues is less than 60%.

10 75. The grain of claim 69 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 12-30 residues is less than 55%.

76. The grain of claim 69 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 31-60 residues is less than about 10%

15 77. The grain of claim 69 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 31-60 residues is less than about 8%

20 78. The grain of claim 77 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 31-60 residues is greater than about 5%,

79. The grain of claim 77 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 31-60 residues is greater than about 6%,

25 80. The grain of claim 37 in a state selected from the group comprising, milled, ground, pearled or rolled, kibbled, cracked, or whole grain.

81. The grain of claim 80 milled to enhance the amount of aleurone layer present.

30 82. The grain of claim 37 having a length to thickness ratio of 4 or less.

83. The grain of claim 37 having a length to thickness ratio of less than about 5.8

84. The grain of claim 37 having a length to thickness ratio of less than about 5.5.

98. The barely plant of claim 89 wherein the starch when gelatinised exhibits reduced peak viscosity.
99. The barley plant of claim 89 wherein the pasting temperature of the starch is elevated.
100. The barley plant of claim 99 wherein the pasting temperature of the starch is higher than 75°C
101. The barley plant of claim 99 wherein the pasting temperature of the starch is higher than 80°C
102. The barley plant of claim 89 wherein the grain has an elevated level of β glucan.
103. The barley plant of claim 102 wherein the β glucan content that is greater than 6% of total non-hulled grain weight
104. The barley plant of claim 102 wherein the β glucan content that is greater than 7% of total non-hulled grain weight
105. The barley plant of claim 102 wherein the β glucan content that is greater than 8% of total non-hulled grain weight
106. The barley plant of claim 102 wherein the β glucan content that is greater than about 15% of total non-hulled grain weight
107. The barley plant of claim 89 wherein the amylose content is higher than 30% (w/w) of the starch content.
108. The barley plant of claim 89 wherein the amylose content is higher than 50% (w/w) of the starch content
109. The barley plant of claim 89 wherein the amylose content is higher than 60% (w/w) of the starch content

122. The barley plant of claim 120 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 6 to 11 residues is greater than 30%.

123. The barley plant of claim 120 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 6 to 11 residues is greater than 35%.

124. The barley plant of claim 120 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 12-30 residues is less than 65%.

125. The barley plant of claim 120 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 12-30 residues is less than 60%.

126. The barley plant of claim 120 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 12-30 residues is less than 55%.

127. The barley plant of claim 120 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 31-60 residues is less than about 10%.

128. The barley plant of claim 120 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 31-60 residues is less than about 8%.

129. The barley plant of claim 128 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 31-60 residues is greater than about 5%.

130. The barley plant of claim 128 wherein the proportion of starch chains that have a degree of polymerisation that falls in the range of 31-60 residues is greater than about 6%.

131. The barley plant of claim 89 in a state selected from the group comprising, milled, ground, pearled or rolled, kibbled, cracked, or whole grain.

132. The barley plant of claim 131 milled to enhance the amount of aleurone layer present.

133. The barley plant of claim 89 having a length to thickness ratio of 4 or less.

134. The barley plant of claim 89 having a length to thickness ratio of less than about 5.8.

SEQUENCE LISTING

<110> Commonwealth Scientific and Industrial Research Organisation
 <120> Barley with reduced SSII activity and starch and starch containing products
 5 with a reduced amylopectin content
 <160> 12
 <210> 1
 <211> 2920
 <212> DNA
 10 <213> Hordeum vulgare
 <220>
 <223> SSII cDNA
 <400> 1
 cctcgagggtg cgtttaccctt acacagagta cactccaact ccagtcgaat ccagcccact 60
 15 gccgcttctg cccgcccctc gtaccgtcgc ccgccccgat cccggccgcc gccatgtcgt 120
 cggcggtcgc gtcccccccg tcccttcctcgc cgctcgcgtc cgccctcgccc gggagatcat 180
 cacggaggag ggcgagggtg ggcgcgtcgc caacccgcgc tggggccggc aggctgcaat 240
 ggcggccgctc gccgctgcag cgcacggctc cgcacggagc ggtggccgcg cgcgcgcgcg 300
 ggatcgacga cgcgcgcgcc ggtaggcagc cccgcgcgtc cgctatggc gccgccacca 360
 20 aggtcgcgga tcccgctcaag acgctcgcgc gcgaagccgc ggaaggtggt gggccgtccc 420
 cgccggcacc gaggcaggac gccgcccgtc tgccgagtaa gaacggcacg ctgatcaacg 480
 gtgagaacaa acctaccggc ggcggtggcg cgactaaaga cagcgggctt gccacaccgc 540
 cacgcgcgcc ccatctgtca atccaaaaca ggtaccggt gaacggtgaa aacaaacata 600
 aggtcgcctc gccgcccacc agcatagtgg atgtcgcgtc tccgggttcc gcagctaaca 660
 25 tttccatcag taacaagggtg ccgctgcctc ttgtccagc caagaagacg ccgccgtcgt 720
 ccgttttccc ggccaagaag acgctgcctc cgtccggctc aaattttgtg tccctcgccct 780
 ctgctcccag gctggacact gtcagcgatg tggaacttgc acagaagaag gatgcgtga 840
 ttgtcaaaaga agctccaaaa ccaaaggctc tttcggcccc tgcaagcccc gctgtacaag 900
 aagacctttg ggatttcaag aaatacattg gtttcgagga gcccggtggg gccaaaggatg 960
 30 atggctcggc tgttcgagat gatgcgggtt cctttgaaca tcaccagaat catgattccg 1020
 gacctttggc aggggagaac gtcagtgatg tggaacttgc tgctgtgaa tgtttccctt 1080
 ggtgcaaaac aggtggtcct ggagatgttg cgggtgcttt gcccaaggct ttggctaaga 1140
 gaggacatcg tgttatgggt gtggtaccaa ggtatgggga ctataggaa gcctacgatg 1200
 tcggagtcgc aaaatactac aaggctgctg gacaggatat ggaagtgaat tatttccatg 1260
 35 ctatatcga tggagtggat ttcgtgttca ttgacgtcc tctcttccga caccgtcagc 1320
 aagacattta tgggggcagc agacaggaaa ttatgaagcg catgattttg ttctgcaagg 1380
 ccgctgtcga ggttcctttg cacgttccat gcggcggtgt cccttacggg gatggaaatc 1440
 tggctctcat tgcaaatgat tggcacacgg cactcctgcc tgtctatctg aaagcatatt 1500
 acagggacca tggtttgatg caatacagtc gctccgttat ggtgatacat aacatcgctc 1560
 40 accaggggccg tggccctgta gatgaattcc gcgttaccca gttgcctgag cactaccttg 1620
 aacacttcag actgtacgac cccgtcggcg gtgagcacgc caactacttc gccgcgggcc 1680
 tgaagatggc ggaccagggt gtgcgtcgtg gccccgggta cctgtgggag ctgaagacgg 1740
 tggaggggcg ctgggggctt cagacatca tacggcagaa cgactggag acccgcgga 1800
 tcgtgaacgg catcgacaac atggagtggg accctgaggt ggacgtccac ctgaagtcgg 1860
 45 acggctacac caacttctcc ctgaagacgc tggactccgg caagcggcag tgcaaggagg 1920
 ccctgcagcg cgagctgggg ctgcaggtcc gcggcgacgt gccgctgctc gggttcatcg 1980
 ggcggctgga cgggcagaag ggcgtggaga tcatcgcgga cgcgatgccc tggatcgtga 2040
 gccaggacgt gcagctgggt atgctgggca cggggcgcca cgacctggag agcatgctgc 2100
 50 agcacttcga gcgggagcac cagacaagg tgcgcgggtg ggtgggggtt tccgtgcgcc 2160
 tggcgccacc gatcacggcg ggcgcgcgac cgctcctcat gccctcccgg ttcgagccgt 2220
 gcgggctgaa ccagctctac gcgatggcct acggcacctt ccccgctcgt caccgcgtcg 2280
 gcggcttgag ggataccgtg ccgcccgttc accccttcaa cactcctggg ctcggttgga 2340
 cgttcgaccg cgcggaggcg cacaagctga tcgaggcgct cgggcaactg ctcgcacct 2400
 accgggacca caaggagagc tggagggggc tccaggagcg cggcatgtcg caggacttca 2460
 55 gctgggaaca tgcgcgaag ctctacgagg acgtcctcgt ccaggccaag taccagtgg 2520
 gaacgctgct accgggtcca gcccgcgat cgtgcattag aggatggaaa tgcgattgc 2580
 gcacttgtag atttggcgca cgcaggaacg tgccgtcctt cttgatgaga acgcccgc 2640
 ccgcgaggtt gagacgtgta ttccgatctg gtccgtcgca gtagtagtg aaacgtcct 2700
 60 tgttgacagt atattgggaat gtttttttct ctttttttct gcgaggagg tatatggga 2760
 tgtaacttg gtattgtaat gtggtatgct gtgtgcatta ttacatcgg tgtgtgtgt 2820
 tattcttgct agctaagtcg gaggccaaag gcgaagcta gctcacatgt ctgatgtatg 2880
 caagtacat ggttggtttg aaaaaaaaaa aaaaaaaaaa 2920
 <210> 2
 <211> 2950

	<212> DNA	
	<213> Hordeum vulgare	
	<220>	
	<221> mutation	
5	<222> 235	
	<223> Barley Line MK6827 cDNA of SSII gene	
	<400> 2	
	gtgcgtttac cccacacaga gtacactcca actccagtc agtccagccc actgccgctt 60	
	ctgccccccc atcgtaccgt cgcccccccc gatccccggc gccgccatgt cgtcggcggt 120	
10	cgcgcccccc gcgtccttcc tcgcgctcgc gtccgcctcg cccgggagat catcacggag 180	
	gagggcgagg gtgggcgcgt cgccaacccg cgctggggcc ggcaggctgc aatgacggcc 240	
	gtcgcgcgtg cagcgcacgg ctgcgcacgg agcgggtggc gcgcgcgcgc cggggatcga 300	
	cgacgcgcgc cccggtaggc agccccgcgc tcgcgcctat ggcgccgcca ccaaggctgc 360	
	ggatcccgtc aagacgctcg atcgcgacgc gcgcgaagg gtggggccgt ccccgccggc 420	
15	accgaggcag gacgcgcgcc gtctgccgag taagaacggc acgctgatca acggtgagaa 480	
	caaacctacc ggcggcggtg gcgcgactaa agacagcggc ctgcccacac ccgcacgcgc 540	
	gccccatctg tcaatccaga acagagtacc ggtgaacggt gaaaacaaac ataaggctcg 600	
	ctcgcgcgtg accagcatag tggatgtcgc gtctccgggt tccgcagcta acatttccat 660	
	cagtaacaag gtgccgccgt ccgttgtccc agccaagaag acgcccggc cgtccgtttt 720	
20	cccggccaaag aaggcgcgcgc cgtcgtccgt tgcctccggc aagaagacgc tgcctcgtc 780	
	cggtcctaat tttgtgtcct cggcctctgc tcccaggctg gacactgtca gcgatgtgga 840	
	acttgacacg aagaaggatg cgtgattgt caaagaagct ccaaaaccaa aggtctcttc 900	
	ggccccgtga gccccgcgtg tacaagaaga cctttgggat ttcaagaaat acattggttt 960	
	cgaggagccc gtggaggcca aggatgatgg ctgggctgtt gcagatgatg cgggttcctt 1020	
25	tgaacatcac cagaatcatg attccggacc tttggcaggg gagaacgtca tgaacgtgg 1080	
	cgctcttgct gctgaatgtt ctccctgggt caaaacaggt ggtcttggag atattgctgg 1140	
	tgttttgcgc aaggctttgg ctaagagagg acatcgtgtt atggttgtgg tacciaagta 1200	
	tggggactat gaggaagcct acgatgtcgg agtccgaaaa tactacaagg ctgctggaca 1260	
	ggatatggaa gtgaattatt tccatgctta tatcgatgga gtggattttg tgttcattga 1320	
30	cgctcctctc ttccgacacc gtcagcaaga catttatggg ggagcagac aggaaattat 1380	
	gaagcgcatg attttgttct gcaaggccgc tgcagaggtt ccttggcacg ttccatgcgg 1440	
	cgggtgtcct tacggggatg gaaatctggt cttcattgca aatgattggc acacggcact 1500	
	cctgcctgtc tatctgaaag catattacag ggaccatggt ttgatgcaat acagtcgctc 1560	
	cgttatgggt atacataaca tcgtccacca gggccgtggc cctgtagatg aattcccgtt 1620	
35	caccgagttg cctgagcact acctggaaca cttcagactg tacgacccc tcggcggtga 1680	
	gcacgccaac tacttcgccc ccggcctgaa gatggcggac cagggttgtc tctgtagccc 1740	
	cgggtacctg tgggagctga agacgggtgga gggcggtcgg gggcttcacg acatcatacg 1800	
	gcagaacgac tgggaagacc gcggcatcgt gaacggcatc gacaacatgg agtggaaacc 1860	
	tgaggtggac gtccacctga agtcggacgc ctacaccaac ttctccctga agacgctgga 1920	
40	ctccggcaag cggcagtgca aggaggccct gcagcgcgag ctggggctgc aggtccgcgg 1980	
	cgacgtgccc ctgctcgggt tcatcgggcg gctggacggg cagaagggcg tggagatcat 2040	
	cgcgacgcgc atgccttgga tcgtgagcca ggacgtgcag ctggtgatgc tgggcacggg 2100	
	gcgccacgac ctggagagca tgcctgagca cttcagcgg gagcaccacg acaagggtgcg 2160	
45	cgggtgggtg ggggtctccg tgcgcctgca gcaccggatc acggcgggcg ccgacgcgtc 2220	
	cctcatgccc tcccgggttc agccgtgcgg gctgaaccag ctctacgcga tggcctacgg 2280	
	caccatccct gtcgtgcacg ccgtcggcgg cctgagggat accgtgcccgc cgttcgaccc 2340	
	cttcaaccac tccgggctcg ggtggacgtt cgaccgcgcc gaggcgcaca agctgatcga 2400	
	ggcgctcggg cactgcctcc gcacctaccg ggaccacaag gagagctgga ggggcctcca 2460	
50	ggagcgcggc atgtcgcagg acttcagctg ggaacatgcc gccaaagtct acgaggacgt 2520	
	cctcgtccag gccaaagtacc agtgggtgaa cctgctaccc ggtccagccc cgcattgcgtg 2580	
	catgagagga tggaaatgca cattgcgcac ttgcagattt ggcgcatgca ggaacgtgcc 2640	
	gtccttcttg atgggaacgc cgcatccgc gcggttgaga cgctgattcc gatctggttc 2700	
	gtcgcagagt agagtgaac gctccttgtt gcaggtatat gggaatgttt tttttttcct 2760	
	tttttttttt gcgagggagg tatatgggaa tgttaacttg gtattgtaat gtggtatgct 2820	
55	gtgtgcatta ttacatcggt tgtgtgtgct tattcttgc agctaagtcg gagccaaga 2880	
	gcgaaagcta gctcacatgt ctgatgtatg caagtgcacat ggttggtttg gttgtgcagt 2940	
	gcaaacggca	2950
	<210> 3	
	<211> 2951	
60	<212> DNA	
	<213> Hordeum vulgare	
	<220>	
	<223> Barley Cultivar Morex cDNA of SSII gene	
	<400> 3	

	gtgcggtttac	cccacacaga	gtacactcca	actccagtc	agtcacagccc	actgccgctt	60
	ctgcccgcgc	atcgtaaccgt	cgcccgcgc	gatcccggcc	gccgccatgt	cgtcggcggt	120
	cgcgcccc	gcgtcccttc	tcgcgctcgc	gtccgcctcg	cccgggagat	catcacggag	180
	gagggcgagg	gtgggcgcg	cgccaaccgc	cgctggggcc	ggcaggctgc	aatggcgcc	240
5	gtcgcgcgtg	cagcgccacg	ctcgcgacgg	agcgggtggc	gcgcgcgcgc	ccgggatcga	300
	cgacgccgcg	cccggtaggc	agccccgcgc	tcgcgcgtat	ggcgcgcga	ccaaggtcgc	360
	ggatcccgtc	aagacgctcg	atcgcgacgc	cgcggaagg	ggtggggcgt	ccccgccggc	420
	accgaggcag	gacggcgccc	gtctgccgag	taagaacggc	acgctgatca	acggtgagaa	480
	caaacctacc	ggcgggcggtg	gcgcgactaa	agacagcggg	ctgcccacac	ccgcacgcgc	540
10	gccccatctg	tcaatccaga	acagagtacc	ggtgaacgg	gaaaaacaa	ataaggtcgc	600
	ctcgcgcgcg	accagcatag	tggatgtcgc	gtctccgggt	tccgcagcta	acatttccat	660
	cagtaacaag	gtgccgcgt	ccgttgtccc	agccaagaag	acgccgcgt	cgctcggttt	720
	cccggccaag	aaggcgccgc	cgctgtccgt	tgtcccggcc	aagaagacgc	tgccgtcgtc	780
	cggtccaaat	tttgtgtcct	cgccctctgc	tcccaggctg	gacactgtca	gcgatgtgga	840
15	acttgacag	aagaaggatg	cgctgattgt	caaagaagct	ccaaaaccaa	aggctctttc	900
	ggcccctgca	gccccgcgt	tacaagaaga	cctttgggat	ttcaagaaat	acattggttt	960
	cgaggagccc	gtggaggcca	aggatgatgg	ctcggctgtt	gcgatgatg	cggttctctt	1020
	tgaacatcac	cagaatcatg	attccggacc	tttgccagg	gagaacgtca	tgaacgtggt	1080
	cgctcggtgt	gctgaatgtt	ctccctgtgt	caaaacaggt	ggtcttgagg	atgttgccgg	1140
20	tgctttgccc	aaggcttttg	ctaagagagg	acatcggtgt	atggtttgtg	taccaaggta	1200
	tggggactat	gaggaagcct	acgatgtcgg	agtcgaaaa	tactacaagg	ctgctggaca	1260
	ggatatggaa	gtgaattatt	tccatgctta	tatcgatgga	gtggattttg	tgctcattga	1320
	cgctccctct	ttccgacacc	gtcagcaaga	catttatggg	ggcagcagac	aggaaattat	1380
	gaagcgcatg	attttgttct	gcaaggccgc	tgctcaggtt	ccttgccacg	ttccatgcgg	1440
25	cggtgtccct	tacggggatg	gaaatctggt	cttcattgca	aatgattggc	acacggcact	1500
	cctgcctgtc	tatctgaaag	catattacag	ggaccatggt	tgatgcaat	acagtcgctc	1560
	cgttatggtg	atacataaca	tcgctcacca	gggcccgtgg	cctgtagatg	aattcccgtt	1620
	caccgagttg	cctgagcact	acctggaaca	cttcagactg	tacgaccccg	tcggcggtga	1680
	gcacgccaac	tacttcgccc	ccggcctgaa	gatggcggac	caggttgtcg	tcgtgagccc	1740
30	cggttacctg	tgggagctga	agacggtgga	ggcgcgctgg	gggcttcacg	acatcatacg	1800
	gcagaacgac	tgggaagacc	gcggcatcgt	gaacggcatc	gacaacatgg	agtggaaacc	1860
	tgaggtggac	gtccacctga	agtcggacgg	ctacaccaac	ttctccctga	agacgctgga	1920
	ctccggcaag	cggcagtgca	aggaggccct	gcagcgcgag	ctggggctgc	aggctccgcg	1980
	cgacgtgcgc	ctgctcgggt	tcacgcggcg	gctggacggg	cagaaggcg	tggagatcat	2040
35	cgcgacgcgc	atgccctgga	tcgtgagcca	ggacgtgcag	ctggtgatgc	tgggcacggg	2100
	gcgccacgac	ctggagagca	tgctgcagca	cttcgagcgg	gagcaccacg	acaaggtgcg	2160
	cggttggtgg	gggttctccg	tgccgctggc	gcaccggatc	acggcgggcg	ccgacgcgct	2220
	cctcatgccc	tcccgggtcg	agccgtgcgg	gctgaaccag	ctctacgcga	tggcctacgg	2280
	caccatccct	gtcgtgcacg	ccgtcggcgg	cctgagggat	accgtgcgcg	cgttcgaccc	2340
40	cttcaaccac	tccgggctcg	ggtggacgtt	cgaccgcgcc	gaggcgcaca	agctgatcga	2400
	ggcgctcggg	cactgcctcc	gcacctaccg	ggaccacaag	gagagctgga	ggggcctcca	2460
	ggagcgcggc	atgctgcagg	acttcagctg	ggaacatgcc	gccaagctct	acgaggacgt	2520
	cctcgtccag	gccaagtacc	agtgggtgaac	gctgctaccc	ggctccagccc	cgcatgcgtg	2580
45	catgagagga	tggaaatgcg	cattgcgcac	ttgcagattt	ggcgcatgca	ggaacgtgcc	2640
	gtccttcttg	atgggaacgc	cggcatccgc	gaggttgaga	cgctgattcc	gatctggctc	2700
	gtcgcagagt	agagtgaac	gtccttgttt	gcaggtatat	gggaatgttt	tttttttcc	2760
	tttttttttt	tgcgaggag	gtatatggga	atgttaactt	ggtattgtaa	tgtggatgac	2820
	tgtgtgcatt	attacatcgg	ttgttgttgc	ttattcttgc	tagctaagtc	ggaggccaag	2880
	agcgaagct	agctcacatg	tctgatgtat	gcaagtgaca	tggttggttt	ggttgtgcag	2940
50	tgcaaacggc	a					2951
	<210>	4					
	<211>	2946					
	<212>	DNA					
	<213>	Hordeum vulgare					
55	<220>						
	<221>	mutation					
	<222>	1855					
	<223>	Barley Line 292 cDNA of SSII gene					
	<400>	4					
60	gtgcggtttac	cccacacaga	gtacactcca	actccagtc	agtcacagccc	actgccgctt	60
	ctgcccgcgc	atcgtaaccgt	cgcccgcgc	gatcccggcc	gccgccatgt	cgtcggcggt	120
	cgcgcccc	gcgtcccttc	tcgcgctcgc	gtccgcctcg	cccgggagat	catcacggag	180
	gagggcgagg	gtgggcgcg	cgccaaccgc	cgctggggcc	ggcaggctgc	aatggcgcc	240
	gtcgcgcgtg	cagcgccacg	ctcgcgacgg	agcgggtggc	gcgcgcgcgc	ccgggatcga	300

	cgacgccg	cccggtaggc	agccccgcgc	tcgcgcgtat	ggcgccgcca	ccaaggtcgc	360
	ggatccccgtc	aagacgcctcg	atcgcgacgc	cgcggaaggt	ggtgggcccgt	ccccgccggc	420
	accgaggcag	gacgccgccc	gtctgcccag	taagaacggc	acgctgatca	acgggtgagaa	480
	caaacctacc	ggcggcggtg	gcgcgactaa	agacagcggg	ctgcccacac	ccgcacgcgc	540
5	gccccatctg	tcaatccaga	acagagtacc	ggtgaacggt	gaaaacaaac	ataaggtcgc	600
	ctcgcgcgcg	accagcatag	tggatgtcgc	gtctccgggt	tccgcagcca	acattttccat	660
	cagtaacaag	gtgccgccc	ccgttgtccc	agccaagaag	acgcgcgcgt	cgtccgtttt	720
	cccggccaa	aaggcgccgc	cgctgcctcg	tgtcccgccc	aagaagacgc	tgccgtcgtc	780
	cggctcaaat	tttgtgtccc	cgccctctgc	tcccaggctg	gacactgtca	gcgatgtgga	840
10	acttgacacag	aagaaggatg	cgctgattgt	caaaagaagct	ccaaaaccaa	aggctctttc	900
	ggccccctgca	gccccgcgtg	tacaagaaga	cccttgggat	ttcaagaaat	acatttggtt	960
	cgaggagccc	gtggaggcca	aggatgatgg	ctcggctggt	gcagatgatg	cgggttcctt	1020
	tgaacatcac	cagaatcatg	attccggacc	tttggcaggg	gagaacgtca	tgaacgtggt	1080
	cgctcgttgc	gctgaatgtt	ctccctggtg	caaaacaggt	ggtcttgagg	atggtgcggg	1140
15	tgctttgccc	aaggctttgg	ctaagagagg	acatcgtggt	atggttggg	taccaaggta	1200
	tggggactat	gaggaagcct	acgatgtcgg	agtcggaaaa	tactacaagg	ctgctggaca	1260
	ggatatggaa	gtgaattatt	tccatgctta	tatcgatgga	gtggattttg	tggtcattga	1320
	gcctcctctc	ttccgacacc	gtcagcaaga	catttatggg	ggcagcagac	aggaaattat	1380
	gaagcgcatg	attttgttct	gcaaggccgc	tgtcggaggt	ccttggcagc	ttccatgcgg	1440
20	cgggtgtccc	tacggggatg	gaaatctggt	cttcattgca	aatgattggc	acacggcact	1500
	cctgcctgtc	tatctgaaag	catattacag	ggaccatggt	ttgatgcaat	acagtcgctc	1560
	cgttatgggtg	atacataaca	tcgctcacca	ggcgcgtggc	cctgtagatg	aattcccgtt	1620
	caccgagttg	cctgagcact	acctggaaca	cttcagactg	tacgaccccg	tcggcggtga	1680
	gcacgccaa	tacttcgccc	ccggcctgaa	gatggcggac	cagggttgtcg	tcgtgagccc	1740
25	cgggtacctg	tgggagctga	agacggtgga	gggcggctgg	gggcttcacg	acatcatacg	1800
	gcagaacgac	tgggaagacc	gcggcatcgt	gaacggcatc	gacaacatgg	agtgaacccc	1860
	tgaggtggac	gtccacctga	agtcggacgg	ctacaccaac	ttctccctga	agacgctgga	1920
	ctccggcaag	cggcagtgca	aggaggccct	gcagcgcgag	ctggggctgc	aggctccgcg	1980
	cgacgtgccc	ctgctcgggt	tcacgggcg	gctggacggg	cagaagggcg	tggagatcat	2040
30	cgcggacgcg	atgccttgga	tcgtgagcca	ggacgtgcag	ctggtgatgc	tgggcacggg	2100
	gcgccacgac	ctggagagca	tgctgcagca	cttcgagcgg	gagcaccacg	acaaggtgcg	2160
	cgggtgggtg	gggtttctccg	tgcgcctggc	gcaccggatc	acggcggggc	ccgacgcgct	2220
	cctcatgccc	tcccgggttcg	agccgtgcgg	gctgaaccag	ctctacgcga	tggcctacgg	2280
	caccatccct	gtcgtgcacg	ccgtcggcgg	cctgagggat	accgtgcgcg	cgttcgaccc	2340
35	cttcaaccac	tccgggctcg	ggtggacgtt	cgaccgcgcc	gaggcgcaca	agctgatcga	2400
	ggcgcctcgg	cactgcctcc	gcacctaccg	ggaccacaag	gagagctgga	ggggcctcca	2460
	ggagcgcggc	atgtcgcagg	acttcagctg	ggaacatgcc	gccaagctct	acgaggacgt	2520
	cctcgtccag	gccaagtacc	agtgggtgaac	gctgctaccc	ggtccagccc	cgcattgcgtg	2580
	catgagagga	tggaaatgcg	cattgcgcac	ttgcagattt	ggcgacgcga	ggaacgtgcc	2640
40	gtcctttctt	atgagaacgc	cggctaccgc	gaggttgaga	cgctgattcc	gatctggtcc	2700
	gtcgcagagt	agagtgaaac	gctccttggt	gcaggtatat	gggaatgttt	tttttccttt	2760
	ttttttgcga	gggaggtata	tgggaatggt	aacttggtat	tgtaatgtgg	tatgctgtgt	2820
	gcattattac	atcgggtggt	gttgcttatt	cttgctagct	aagtcggagg	ccaagagcga	2880
	aagctagctc	acatgtctga	tgatatgcaag	tgacatgggt	ggtttggttg	tgcagtgcga	2940
45	acggca						2946
	<210>	5					
	<211>	813					
	<212>	PRT					
	<213>	Hordeum vulgare					
50	<220>						
	<223>	Barley Cultivar Morex SSII deduced amino acid sequence					
	<400>	5					
	Met Ser Ser Ala Val Ala Ser Pro Ala Ser Phe Leu Ala Leu Ala						
	1 5 10 15						
55	Ser Ala Ser Pro Gly Arg Ser Ser Arg Arg Arg Ala Arg Val Gly						
	20 25 30						
	Ala Ser Pro Thr Arg Ala Gly Ala Gly Arg Leu Gln Trp Arg Pro						
	35 40 45						
	Ser Pro Leu Gln Arg Thr Ala Arg Asp Gly Ala Val Ala Ala Arg						
60	50 55 60						
	Ala Ala Gly Ile Asp Asp Ala Ala Pro Gly Arg Gln Pro Arg Ala						
	65 70 75						
	Arg Arg Tyr Gly Ala Ala Thr Lys Val Ala Asp Pro Val Lys Thr						
	80 85 90						

	Leu	Asp	Arg	Asp	Ala	Ala	Glu	Gly	Gly	Gly	Pro	Ser	Pro	Pro	Ala	
					95					100					105	
	Pro	Arg	Gln	Asp	Ala	Ala	Arg	Leu	Pro	Ser	Lys	Asn	Gly	Thr	Leu	
					110					115					120	
5	Ile	Asn	Gly	Glu	Asn	Lys	Pro	Thr	Gly	Gly	Gly	Gly	Ala	Thr	Lys	
					125					130					135	
	Asp	Ser	Gly	Leu	Pro	Thr	Pro	Ala	Arg	Ala	Pro	His	Leu	Ser	Ile	
					140					145					150	
	Gln	Asn	Arg	Val	Pro	Val	Asn	Gly	Glu	Asn	Lys	His	Lys	Val	Ala	
10					155					160					165	
	Ser	Pro	Pro	Thr	Ser	Ile	Val	Asp	Val	Ala	Ser	Pro	Gly	Ser	Ala	
					170					175					180	
	Ala	Asn	Ile	Ser	Ile	Ser	Asn	Lys	Val	Pro	Pro	Ser	Val	Val	Pro	
					185					190					195	
15	Ala	Lys	Lys	Thr	Pro	Pro	Ser	Ser	Val	Phe	Pro	Ala	Lys	Lys	Ala	
					200					205					210	
	Pro	Pro	Ser	Ser	Val	Val	Pro	Ala	Lys	Lys	Thr	Leu	Pro	Ser	Ser	
					215					220					225	
20	Gly	Ser	Asn	Phe	Val	Ser	Ser	Ala	Ser	Ala	Pro	Arg	Leu	Asp	Thr	
					230					235					240	
	Val	Ser	Asp	Val	Glu	Leu	Ala	Gln	Lys	Lys	Asp	Ala	Leu	Ile	Val	
					245					250					255	
	Lys	Glu	Ala	Pro	Lys	Pro	Lys	Ala	Leu	Ser	Ala	Pro	Ala	Ala	Pro	
					260					265					270	
25	Ala	Val	Gln	Glu	Asp	Leu	Trp	Asp	Phe	Lys	Lys	Tyr	Ile	Gly	Phe	
					275					280					285	
	Glu	Glu	Pro	Val	Glu	Ala	Lys	Asp	Asp	Gly	Ser	Ala	Val	Ala	Asp	
					290					295					300	
30	Asp	Ala	Gly	Ser	Phe	Glu	His	His	Gln	Asn	His	Asp	Ser	Gly	Pro	
					305					310					315	
	Leu	Ala	Gly	Glu	Asn	Val	Met	Asn	Val	Val	Val	Val	Ala	Ala	Glu	
					320					325					330	
	Cys	Ser	Pro	Trp	Cys	Lys	Thr	Gly	Gly	Leu	Gly	Asp	Val	Ala	Gly	
					335					340					345	
35	Ala	Leu	Pro	Lys	Ala	Leu	Ala	Lys	Arg	Gly	His	Arg	Val	Met	Val	
					350					355					360	
	Val	Val	Pro	Arg	Tyr	Gly	Asp	Tyr	Glu	Glu	Ala	Tyr	Asp	Val	Gly	
					365					370					375	
	Val	Arg	Lys	Tyr	Tyr	Lys	Ala	Ala	Gly	Gln	Asp	Met	Glu	Val	Asn	
40					380					385					390	
	Tyr	Phe	His	Ala	Tyr	Ile	Asp	Gly	Val	Asp	Phe	Val	Phe	Ile	Asp	
					395					400					405	
	Ala	Pro	Leu	Phe	Arg	His	Arg	Gln	Gln	Asp	Ile	Tyr	Gly	Gly	Ser	
					410					415					420	
45	Arg	Gln	Glu	Ile	Met	Lys	Arg	Met	Ile	Leu	Phe	Cys	Lys	Ala	Ala	
					425					430					435	
	Val	Glu	Val	Pro	Trp	His	Val	Pro	Cys	Gly	Gly	Val	Pro	Tyr	Gly	
					440					445					450	
	Asp	Gly	Asn	Leu	Val	Phe	Ile	Ala	Asn	Asp	Trp	His	Thr	Ala	Leu	
50					455					460					465	
	Leu	Pro	Val	Tyr	Leu	Lys	Ala	Tyr	Tyr	Arg	Asp	His	Gly	Leu	Met	
					470					475					480	
	Gln	Tyr	Ser	Arg	Ser	Val	Met	Val	Ile	His	Asn	Ile	Ala	His	Gln	
					485					490					495	
55	Gly	Arg	Gly	Pro	Val	Asp	Glu	Phe	Pro	Phe	Thr	Glu	Leu	Pro	Glu	
					500					505					510	
	His	Tyr	Leu	Glu	His	Phe	Arg	Leu	Tyr	Asp	Pro	Val	Gly	Gly	Glu	
					515					520					525	
	His	Ala	Asn	Tyr	Phe	Ala	Ala	Gly	Leu	Lys	Met	Ala	Asp	Gln	Val	
60					530					535					540	
	Val	Val	Val	Ser	Pro	Gly	Tyr	Leu	Trp	Glu	Leu	Lys	Thr	Val	Glu	
					545					550					555	
	Gly	Gly	Trp	Gly	Leu	His	Asp	Ile	Ile	Arg	Gln	Asn	Asp	Trp	Lys	
					560					565					570	

Thr Arg Gly Ile Val Asn Gly Ile Asp Asn Met Glu Trp Asn Pro
 575 580 585
 Glu Val Asp Val His Leu Lys Ser Asp Gly Tyr Thr Asn Phe Ser
 590 595 600
 5 Leu Lys Thr Leu Asp Ser Gly Lys Arg Gln Cys Lys Glu Ala Leu
 605 610 615
 Gln Arg Glu Leu Gly Leu Gln Val Arg Gly Asp Val Pro Leu Leu
 620 625 630
 Gly Phe Ile Gly Arg Leu Asp Gly Gln Lys Gly Val Glu Ile Ile
 635 640 645
 10 Ala Asp Ala Met Pro Trp Ile Val Ser Gln Asp Val Gln Leu Val
 650 655 660
 Met Leu Gly Thr Gly Arg His Asp Leu Glu Ser Met Leu Gln His
 665 670 675
 15 Phe Glu Arg Glu His His Asp Lys Val Arg Gly Trp Val Gly Phe
 680 685 690
 Ser Val Arg Leu Ala His Arg Ile Thr Ala Gly Ala Asp Ala Leu
 695 700 705
 20 Leu Met Pro Ser Arg Phe Glu Pro Cys Gly Leu Asn Gln Leu Tyr
 710 715 720
 Ala Met Ala Tyr Gly Thr Ile Pro Val Val His Ala Val Gly Gly
 725 730 735
 Leu Arg Asp Thr Val Pro Pro Phe Asp Pro Phe Asn His Ser Gly
 740 745 750
 25 Leu Gly Trp Thr Phe Asp Arg Ala Glu Ala His Lys Leu Ile Glu
 755 760 765
 Ala Leu Gly His Cys Leu Arg Thr Tyr Arg Asp His Lys Glu Ser
 770 775 780
 Trp Arg Gly Leu Gln Glu Arg Gly Met Ser Gln Asp Phe Ser Trp
 785 790 795
 30 Glu His Ala Ala Lys Leu Tyr Glu Asp Val Leu Val Gln Ala Lys
 800 805 810
 Tyr Gln Trp ***
 <210> 6
 35 <211> 802
 <212> PRT
 <213> Hordeum vulgare
 <220>
 <223> Barley Cultivar Himalaya SSII deduced amino acid sequence
 40 <400> 6
 Met Ser Ser Ala Val Ala Ser Pro Ala Ser Phe Leu Ala Leu Ala
 1 5 10 15
 Ser Ala Ser Pro Gly Arg Ser Ser Arg Arg Arg Ala Arg Val Gly
 20 25 30
 45 Ala Ser Pro Thr Arg Ala Gly Ala Gly Arg Leu Gln Trp Arg Pro
 35 40 45
 Ser Pro Leu Gln Arg Thr Ala Arg Asp Gly Ala Val Ala Ala Arg
 50 55 60
 Ala Ala Gly Ile Asp Asp Ala Ala Pro Gly Arg Gln Pro Arg Ala
 65 70 75
 50 Arg Arg Tyr Gly Ala Ala Thr Lys Val Ala Asp Pro Val Lys Thr
 80 85 90
 Leu Asp Arg Asp Ala Ala Glu Gly Gly Gly Pro Ser Pro Pro Ala
 95 100 105
 55 Pro Arg Gln Asp Ala Ala Arg Leu Pro Ser Lys Asn Gly Thr Leu
 110 115 120
 Ile Asn Gly Glu Asn Lys Pro Thr Gly Gly Gly Ala Thr Lys
 125 130 135
 Asp Ser Gly Leu Pro Thr Pro Ala Arg Ala Pro His Leu Ser Ile
 140 145 150
 60 Gln Asn Arg Val Pro Val Asn Gly Glu Asn Lys His Lys Val Ala
 155 160 165
 Ser Pro Pro Thr Ser Ile Val Asp Val Ala Ser Pro Gly Ser Ala
 170 175 180

	Ala Asn Ile Ser Ile Ser Asn Lys Val Pro Pro Ser Val Val Pro	185	190	195
	Ala Lys Lys Thr Pro Pro Ser Ser Val Phe Pro Ala Lys Lys Thr	200	205	210
5	Leu Pro Ser Ser Gly Ser Asn Phe Val Ser Ser Ala Ser Ala Pro	215	220	225
	Arg Leu Asp Thr Val Ser Asp Val Glu Leu Ala Gln Lys Lys Asp	230	235	240
	Ala Leu Ile Val Lys Glu Ala Pro Lys Pro Lys Ala Leu Ser Ala	245	250	255
10	Pro Ala Ala Pro Ala Val Gln Glu Asp Leu Trp Asp Phe Lys Lys	260	265	270
	Tyr Ile Gly Phe Glu Glu Pro Val Glu Ala Lys Asp Asp Gly Ser	275	280	285
15	Ala Val Ala Asp Asp Ala Gly Ser Phe Glu His His Gln Asn His	290	295	300
	Asp Ser Gly Pro Leu Ala Gly Glu Asn Val Met Asn Val Val Val	305	310	315
20	Val Ala Ala Glu Cys Ser Pro Trp Cys Lys Thr Gly Gly Leu Gly	320	325	330
	Asp Val Ala Gly Ala Leu Pro Lys Ala Leu Ala Lys Arg Gly His	335	340	345
	Arg Val Met Val Val Val Pro Arg Tyr Gly Asp Tyr Glu Glu Ala	350	355	360
25	Tyr Asp Val Gly Val Arg Lys Tyr Tyr Lys Ala Ala Gly Gln Asp	365	370	375
	Met Glu Val Asn Tyr Phe His Ala Tyr Ile Asp Gly Val Asp Phe	380	385	390
30	Val Phe Ile Asp Ala Pro Leu Phe Arg His Arg Gln Gln Asp Ile	395	400	405
	Tyr Gly Gly Ser Arg Gln Glu Ile Met Lys Arg Met Ile Leu Phe	410	415	420
	Cys Lys Ala Ala Val Glu Val Pro Trp His Val Pro Cys Gly Gly	425	430	435
35	Val Pro Tyr Gly Asp Gly Asn Leu Val Phe Ile Ala Asn Asp Trp	440	445	450
	His Thr Ala Leu Leu Pro Val Tyr Leu Lys Ala Tyr Tyr Arg Asp	455	460	465
40	His Gly Leu Met Gln Tyr Ser Arg Ser Val Met Val Ile His Asn	470	475	480
	Ile Ala His Gln Gly Arg Gly Pro Val Asp Glu Phe Pro Phe Thr	485	490	495
	Glu Leu Pro Glu His Tyr Leu Glu His Phe Arg Leu Tyr Asp Pro	500	505	510
45	Val Gly Gly Glu His Ala Asn Tyr Phe Ala Ala Gly Leu Lys Met	515	520	525
	Ala Asp Gln Val Val Val Val Ser Pro Gly Tyr Leu Trp Glu Leu	530	535	540
50	Lys Thr Val Glu Gly Gly Trp Gly Leu His Asp Ile Ile Arg Gln	545	550	555
	Asn Asp Trp Lys Thr Arg Gly Ile Val Asn Gly Ile Asp Asn Met	560	565	570
	Glu Trp Asn Pro Glu Val Asp Val His Leu Lys Ser Asp Gly Tyr	575	580	585
55	Thr Asn Phe Ser Leu Lys Thr Leu Asp Ser Gly Lys Arg Gln Cys	590	595	600
	Lys Glu Ala Leu Gln Arg Glu Leu Gly Leu Gln Val Arg Gly Asp	605	610	615
60	Val Pro Leu Leu Gly Phe Ile Gly Arg Leu Asp Gly Gln Lys Gly	620	625	630
	Val Glu Ile Ile Ala Asp Ala Met Pro Trp Ile Val Ser Gln Asp	635	640	645
	Val Gln Leu Val Met Leu Gly Thr Gly Arg His Asp Leu Glu Ser	650	655	660


Met Leu Gln His Phe Glu Arg Glu His His Asp Lys Val Arg Gly
 665 670 675
 Trp Val Gly Phe Ser Val Arg Leu Ala His Arg Ile Thr Ala Gly
 680 685 690
 5 Ala Asp Ala Leu Leu Met Pro Ser Arg Phe Glu Pro Cys Gly Leu
 695 700 705
 Asn Gln Leu Tyr Ala Met Ala Tyr Gly Thr Ile Pro Val Val His
 710 715 720
 10 Ala Val Gly Gly Leu Arg Asp Thr Val Pro Pro Phe Asp Pro Phe
 725 730 735
 Asn His Ser Gly Leu Gly Trp Thr Phe Asp Arg Ala Glu Ala His
 740 745 750
 Lys Leu Ile Glu Ala Leu Gly His Cys Leu Arg Thr Tyr Arg Asp
 755 760 765
 15 His Lys Glu Ser Trp Arg Gly Leu Gln Glu Arg Gly Met Ser Gln
 770 775 780
 Asp Phe Ser Trp Glu His Ala Ala Lys Leu Tyr Glu Asp Val Leu
 785 790 795
 Val Gln Ala Lys Tyr Gln Trp ***
 800
 20
 <210> 7
 <211> 813
 <212> PRT
 <213> Hordeum vulgare
 25
 <220>
 <221> SITE
 <222> 583
 <223> Barley Line 292 SSII deduced amino acid sequence with premature termination
 <400> 7
 30 Met Ser Ser Ala Val Ala Ser Pro Ala Ser Phe Leu Ala Leu Ala
 1 5 10 15
 Ser Ala Ser Pro Gly Arg Ser Ser Arg Arg Arg Ala Arg Val Gly
 20 25 30
 35 Ala Ser Pro Thr Arg Ala Gly Ala Gly Arg Leu Gln Trp Arg Pro
 35 40 45
 Ser Pro Leu Gln Arg Thr Ala Arg Asp Gly Ala Val Ala Ala Arg
 50 55 60
 Ala Ala Gly Ile Asp Asp Ala Ala Pro Gly Arg Gln Pro Arg Ala
 65 70 75
 40 Arg Arg Tyr Gly Ala Ala Thr Lys Val Ala Asp Pro Val Lys Thr
 80 85 90
 Leu Asp Arg Asp Ala Ala Glu Gly Gly Gly Pro Ser Pro Pro Ala
 95 100 105
 45 Pro Arg Gln Asp Ala Ala Arg Leu Pro Ser Lys Asn Gly Thr Leu
 110 115 120
 Ile Asn Gly Glu Asn Lys Pro Thr Gly Gly Gly Gly Ala Thr Lys
 125 130 135
 Asp Ser Gly Leu Pro Thr Pro Ala Arg Ala Pro His Leu Ser Ile
 140 145 150
 50 Gln Asn Arg Val Pro Val Asn Gly Glu Asn Lys His Lys Val Ala
 155 160 165
 Ser Pro Pro Thr Ser Ile Val Asp Val Ala Ser Pro Gly Ser Ala
 170 175 180
 55 Ala Asn Ile Ser Ile Ser Asn Lys Val Pro Pro Ser Val Val Pro
 185 190 195
 Ala Lys Lys Thr Pro Pro Ser Ser Val Phe Pro Ala Lys Lys Ala
 200 205 210
 Pro Pro Ser Ser Val Val Pro Ala Lys Lys Thr Leu Pro Ser Ser
 215 220 225
 60 Gly Ser Asn Phe Val Ser Ser Ala Ser Ala Pro Arg Leu Asp Thr
 230 235 240
 Val Ser Asp Val Glu Leu Ala Gln Lys Lys Asp Ala Leu Ile Val
 245 250 255
 Lys Glu Ala Pro Lys Pro Lys Ala Leu Ser Ala Pro Ala Ala Pro

		260		265		270
		Ala Val Gln Glu Asp Leu Trp Asp Phe Lys Lys Tyr Ile Gly Phe				
		275		280		285
5		Glu Glu Pro Val Glu Ala Lys Asp Asp Gly Ser Ala Val Ala Asp				
		290		295		300
		Asp Ala Gly Ser Phe Glu His His Gln Asn His Asp Ser Gly Pro				
		305		310		315
		Leu Ala Gly Glu Asn Val Met Asn Val Val Val Ala Ala Glu				
		320		325		330
10		Cys Ser Pro Trp Cys Lys Thr Gly Gly Leu Gly Asp Val Ala Gly				
		335		340		345
		Ala Leu Pro Lys Ala Leu Ala Lys Arg Gly His Arg Val Met Val				
		350		355		360
		Val Val Pro Arg Tyr Gly Asp Tyr Glu Glu Ala Tyr Asp Val Gly				
15		365		370		375
		Val Arg Lys Tyr Tyr Lys Ala Ala Gly Gln Asp Met Glu Val Asn				
		380		385		390
		Tyr Phe His Ala Tyr Ile Asp Gly Val Asp Phe Val Phe Ile Asp				
		395		400		405
20		Ala Pro Leu Phe Arg His Arg Gln Gln Asp Ile Tyr Gly Gly Ser				
		410		415		420
		Arg Gln Glu Ile Met Lys Arg Met Ile Leu Phe Cys Lys Ala Ala				
		425		430		435
		Val Glu Val Pro Trp His Val Pro Cys Gly Gly Val Pro Tyr Gly				
25		440		445		450
		Asp Gly Asn Leu Val Phe Ile Ala Asn Asp Trp His Thr Ala Leu				
		455		460		465
		Leu Pro Val Tyr Leu Lys Ala Tyr Tyr Arg Asp His Gly Leu Met				
		470		475		480
30		Gln Tyr Ser Arg Ser Val Met Val Ile His Asn Ile Ala His Gln				
		485		490		495
		Gly Arg Gly Pro Val Asp Glu Phe Pro Phe Thr Glu Leu Pro Glu				
		500		505		510
		His Tyr Leu Glu His Phe Arg Leu Tyr Asp Pro Val Gly Gly Glu				
35		515		520		525
		His Ala Asn Tyr Phe Ala Ala Gly Leu Lys Met Ala Asp Gln Val				
		530		535		540
		Val Val Val Ser Pro Gly Tyr Leu Trp Glu Leu Lys Thr Val Glu				
		545		550		555
40		Gly Gly Trp Gly Leu His Asp Ile Ile Arg Gln Asn Asp Trp Lys				
		560		565		570
		Thr Arg Gly Ile Val Asn Gly Ile Asp Asn Met Glu *** Asn Pro				
		575		580		585
		Glu Val Asp Val His Leu Lys Ser Asp Gly Tyr Thr Asn Phe Ser				
45		590		595		600
		Leu Lys Thr Leu Asp Ser Gly Lys Arg Gln Cys Lys Glu Ala Leu				
		605		610		615
		Gln Arg Glu Leu Gly Leu Gln Val Arg Gly Asp Val Pro Leu Leu				
		620		625		630
50		Gly Phe Ile Gly Arg Leu Asp Gly Gln Lys Gly Val Glu Ile Ile				
		635		640		645
		Ala Asp Ala Met Pro Trp Ile Val Ser Gln Asp Val Gln Leu Val				
		650		655		660
		Met Leu Gly Thr Gly Arg His Asp Leu Glu Ser Met Leu Gln His				
55		665		670		675
		Phe Glu Arg Glu His His Asp Lys Val Arg Gly Trp Val Gly Phe				
		680		685		690
		Ser Val Arg Leu Ala His Arg Ile Thr Ala Gly Ala Asp Ala Leu				
		695		700		705
60		Leu Met Pro Ser Arg Phe Glu Pro Cys Gly Leu Asn Gln Leu Tyr				
		710		715		720
		Ala Met Ala Tyr Gly Thr Ile Pro Val Val His Ala Val Gly Gly				
		725		730		735
		Leu Arg Asp Thr Val Pro Pro Phe Asp Pro Phe Asn His Ser Gly				

740 745 750
 Leu Gly Trp Thr Phe Asp Arg Ala Glu Ala His Lys Leu Ile Glu
 755 760 765
 5 Ala Leu Gly His Cys Leu Arg Thr Tyr Arg Asp His Lys Glu Ser
 770 775 780
 Trp Arg Gly Leu Gln Glu Arg Gly Met Ser Gln Asp Phe Ser Trp
 785 790 795
 Glu His Ala Ala Lys Leu Tyr Glu Asp Val Leu Val Gln Ala Lys
 800 805 810
 10 Tyr Gln Trp ***
 <210> 8
 <211> 813
 <212> PRT
 <213> Hordeum vulgare
 15 <220>
 <221> SITE
 <222> 43
 <223> Barley Line MK5827 SSII deduced amino acid sequence with premature
 termination
 20 <400> 8
 Met Ser Ser Ala Val Ala Ser Pro Ala Ser Phe Leu Ala Leu Ala
 1 5 10 15
 Ser Ala Ser Pro Gly Arg Ser Ser Arg Arg Arg Ala Arg Val Gly
 20 25 30
 25 Ala Ser Pro Thr Arg Ala Gly Ala Gly Arg Leu Gln *** Arg Pro
 35 40 45
 Ser Pro Leu Gln Arg Thr Ala Arg Asp Gly Ala Val Ala Ala Arg
 50 55 60
 Ala Ala Gly Ile Asp Asp Ala Ala Pro Gly Arg Gln Pro Arg Ala
 30 65 70 75
 Arg Arg Tyr Gly Ala Ala Thr Lys Val Ala Asp Pro Val Lys Thr
 80 85 90
 Leu Asp Arg Asp Ala Ala Glu Gly Gly Gly Pro Ser Pro Pro Ala
 95 100 105
 35 Pro Arg Gln Asp Ala Ala Arg Leu Pro Ser Lys Asn Gly Thr Leu
 110 115 120
 Ile Asn Gly Glu Asn Lys Pro Thr Gly Gly Gly Gly Ala Thr Lys
 125 130 135
 Asp Ser Gly Leu Pro Thr Pro Ala Arg Ala Pro His Leu Ser Ile
 40 140 145 150
 Gln Asn Arg Val Pro Val Asn Gly Glu Asn Lys His Lys Val Ala
 155 160 165
 Ser Pro Pro Thr Ser Ile Val Asp Val Ala Ser Pro Gly Ser Ala
 170 175 180
 45 Ala Asn Ile Ser Ile Ser Asn Lys Val Pro Pro Ser Val Val Pro
 185 190 195
 Ala Lys Lys Thr Pro Pro Ser Ser Val Phe Pro Ala Lys Lys Ala
 200 205 210
 Pro Pro Ser Ser Val Val Pro Ala Lys Lys Thr Leu Pro Ser Ser
 50 215 220 225
 Gly Ser Asn Phe Val Ser Ser Ala Ser Ala Pro Arg Leu Asp Thr
 230 235 240
 Val Ser Asp Val Glu Leu Ala Gln Lys Lys Asp Ala Leu Ile Val
 245 250 255
 55 Lys Glu Ala Pro Lys Pro Lys Ala Leu Ser Ala Pro Ala Ala Pro
 260 265 270
 Ala Val Gln Glu Asp Leu Trp Asp Phe Lys Lys Tyr Ile Gly Phe
 275 280 285
 Glu Glu Pro Val Glu Ala Lys Asp Asp Gly Ser Ala Val Ala Asp
 290 295 300
 60 Asp Ala Gly Ser Phe Glu His His Gln Asn His Asp Ser Gly Pro
 305 310 315
 Leu Ala Gly Glu Asn Val Met Asn Val Val Val Val Ala Ala Glu
 320 325 330

	Cys Ser Pro Trp Cys Lys Thr Gly Gly Leu Gly Asp Ile Ala Gly		
	335	340	345
	Ala Leu Pro Lys Ala Leu Ala Lys Arg Gly His Arg Val Met Val		
	350	355	360
5	Val Val Pro Arg Tyr Gly Asp Tyr Glu Glu Ala Tyr Asp Val Gly		
	365	370	375
	Val Arg Lys Tyr Tyr Lys Ala Ala Gly Gln Asp Met Glu Val Asn		
	380	385	390
	Tyr Phe His Ala Tyr Ile Asp Gly Val Asp Phe Val Phe Ile Asp		
10	395	400	405
	Ala Pro Leu Phe Arg His Arg Gln Gln Asp Ile Tyr Gly Gly Ser		
	410	415	420
	Arg Gln Glu Ile Met Lys Arg Met Ile Leu Phe Cys Lys Ala Ala		
	425	430	435
15	Val Glu Val Pro Trp His Val Pro Cys Gly Gly Val Pro Tyr Gly		
	440	445	450
	Asp Gly Asn Leu Val Phe Ile Ala Asn Asp Trp His Thr Ala Leu		
	455	460	465
	Leu Pro Val Tyr Leu Lys Ala Tyr Tyr Arg Asp His Gly Leu Met		
20	470	475	480
	Gln Tyr Ser Arg Ser Val Met Val Ile His Asn Ile Ala His Gln		
	485	490	495
	Gly Arg Gly Pro Val Asp Glu Phe Pro Phe Thr Glu Leu Pro Glu		
	500	505	510
25	His Tyr Leu Glu His Phe Arg Leu Tyr Asp Pro Val Gly Gly Glu		
	515	520	525
	His Ala Asn Tyr Phe Ala Ala Gly Leu Lys Met Ala Asp Gln Val		
	530	535	540
	Val Val Val Ser Pro Gly Tyr Leu Trp Glu Leu Lys Thr Val Glu		
30	545	550	555
	Gly Gly Trp Gly Leu His Asp Ile Ile Arg Gln Asn Asp Trp Lys		
	560	565	570
	Thr Arg Gly Ile Val Asn Gly Ile Asp Asn Met Glu Trp Asn Pro		
	575	580	585
35	Glu Val Asp Val His Leu Lys Ser Asp Gly Tyr Thr Asn Phe Ser		
	590	595	600
	Leu Lys Thr Leu Asp Ser Gly Lys Arg Gln Cys Lys Glu Ala Leu		
	605	610	615
	Gln Arg Glu Leu Gly Leu Gln Val Arg Gly Asp Val Pro Leu Leu		
40	620	625	630
	Gly Phe Ile Gly Arg Leu Asp Gly Gln Lys Gly Val Glu Ile Ile		
	635	640	645
	Ala Asp Ala Met Pro Trp Ile Val Ser Gln Asp Val Gln Leu Val		
	650	655	660
45	Met Leu Gly Thr Gly Arg His Asp Leu Glu Ser Met Leu Gln His		
	665	670	675
	Phe Glu Arg Glu His His Asp Lys Val Arg Gly Trp Val Gly Phe		
	680	685	690
	Ser Val Arg Leu Ala His Arg Ile Thr Ala Gly Ala Asp Ala Leu		
50	695	700	705
	Leu Met Pro Ser Arg Phe Glu Pro Cys Gly Leu Asn Gln Leu Tyr		
	710	715	720
	Ala Met Ala Tyr Gly Thr Ile Pro Val Val His Ala Val Gly Gly		
	725	730	735
55	Leu Arg Asp Thr Val Pro Pro Phe Asp Pro Phe Asn His Ser Gly		
	740	745	750
	Leu Gly Trp Thr Phe Asp Arg Ala Glu Ala His Lys Leu Ile Glu		
	755	760	765
	Ala Leu Gly His Cys Leu Arg Thr Tyr Arg Asp His Lys Glu Ser		
60	770	775	780
	Trp Arg Gly Leu Gln Glu Arg Gly Met Ser Gln Asp Phe Ser Trp		
	785	790	795
	Glu His Ala Ala Lys Leu Tyr Glu Asp Val Leu Val Gln Ala Lys		
	800	805	810

Tyr Gln Trp ***
<210> 9
<211> 23
<212> DNA
5 <213> Artificial
<220>
<223> PCR primer ssIIa
<400> 9
tggttgagggtt ccatggcacg ttc 23
10 <210> 10
<211> 23
<212> DNA
<213> Artificial
<220>
15 <223> PCR primer ssIIb
<400> 10
agtcgttctg ccgtatgatg tcg 23
20 <210> 11
<211> 23
<212> DNA
<213> Artificial
<220>
<223> PCR primer ZLSS2P4
<400> 11
25 cctggaacac ttcagactgt acg 23
<210> 12
<211> 23
<212> DNA
<213> Artificial
30 <220>
<223> PCR primer ZLBSSII5
<400> 12
cttcaggag aagttggtgt agc 23
35



2/23




FIGURE 2

3/23




FIGURE 3

4/23




FIGURE 4




FIGURE 5

6/23




FIGURE 6

7/23

70	Rpg1 (T)	Reaction to <i>Puccinia graminis</i> 1
67	Run1	Reaction to <i>Ustilago nuda</i> 1
61	brh1 (br, ari-i)	Brachytic 1 (Breviaristatum-i)
52	fch12 (fc)	Chlorina seedling 12 (Chlorina seedling c)
50	wax (wx)	Waxy endosperm
48	gsh3 (cer-a) Glossy sheath 3 (Eceriferum-a)	- - ++
43	fch5 (f5)	Chlorina seedling 5
39	yvs2 (yc)	Virescent seedling 2 (Yellow seedling c)
36	cer-ze (gl5)	Eceriferum-ze (Glossy leaf 5) ++ ++ -
32	wnd	Winding dwarf
	rsm1 (sm)	Reaction to BSMV 1
26	abo7 (a ω)	Albino seedling 7 (Albino seedling c2)
23	ant1 (Rs)	Anthocyanin-less 1 (Red stem)
22	ert-m	Erectoides-m
18	ert-a	Erectoides-a
13	ert-d	Erectoides-d
12	fch8 (f8)	Chlorina seedling 8
10	fst3 (fs3)	Fragile stem 3
9	cer-f	Eceriferum-f + + ++
8	dsp1 (l)	Dense spike 1
7	msg14	Male sterile genetic 14
6	msg10	Male sterile genetic 10
4	sex6	Shrunken endosperm xenia 6
2	seg5	Shrunken endosperm 5
1	seg2	Shrunken endosperm 2
0	nud (n)	Naked caryopsis
- 5	fch4 (f4)	Chlorina seedling 4
- 6	Amy2	Alpha-amylase 2
- 7	lks2 (lk2)	Short awn 2
- 8	ubs4 (u4, ari-d)	Unbranched style 4 (Breviaristatum-d)
- 11	blx2 (bl2)	Non-blue aleurone xenia 2
- 20	lbi3 (lb3)	Long basal rachis internode 3
	xnt4 (x ω)	Xantha seedling 4 (Xantha seedling c2)
	msg50	Male sterile genetic 50
- 31	Rym2 (Ym2)	Reaction to BaYMV 2
- 35	seg4	Shrunken endosperm 4
- 46	Xnt1 (X α)	Xantha seedling 1 (Xantha a)
- 56	Rph3 (Pa3) Reaction to <i>Puccinia hordei</i> 3	
	xnt9 (xan, i)	Xantha seedling 9 (Xantha seedling i)
- 80	seg1	Shrunken endosperm 1
- 85	msg23	Male sterile genetic 23

FIGURE 7

8/23





FIGURE 8



9/23

Barley SSII cDNA Sequence

FIGURE 9

```

1  CCTCGAGGTG CGTTTACCCC ACACAGAGTA CACTCCAACCT CCAGTCCAAT
51  CCAGCCCACT GCCGCTTCTG CCCGCCCATC GTACCGTCGC CCGCCCCGAT
101 CCCGGCCGCC GCCATGTCGT CGGCGGTCGC GTCCCCCGCG TCCTTCCTCG
151 CGCTCGCGTC CGCCTCGCCC GGGAGATCAT CACGAGGAG GGCAGGGGTG
201 GGCCTGTCGC CAACCCGCGC TGGGGCCGGC AGGCTGCAAT GCGGCCCGTC
251 GCCGCTGCAG CGCACGGCTC GCGACGGAGC GGTGGCCGCG CCGCCCGCCG
301 GGATCGACGA CGCCGCGCCC GGTAGGCAGC CCCGCGCTCG CCGCTATGGC
351 GCCGCCACCA AGGTCGCGGA TCCCCTCAAG ACGCTCGATC GCGACGCCGC
401 GGAAGGTGGT GGGCCGTCCC CGCCGGCACC GAGGCAGGAC GCCGCCCGTC
451 TGCCGAGTAA GAACGGCACG CTGATCAACG GTGAGAACA ACCTACCGGC
501 GGCCTGGCG CGACTAAAGA CAGCGGGCTT GCCACACCCG CACGCGCGCC
551 CCATCTGTCA ATCCAAACA GAGTACCGGT GAACGGTGAA AACAAACATA
601 AGGTGCGCTC GCCGCCGACC AGCATAGTGG ATGTCGCGTC TCCGGGTTC
651 GCAGCTAACA TTTCCATCAG TAACAAGGTG CCGCCGTCCG TTGTCCCAGC
701 CAAGAAGACG CCGCCGTCGT CCGTTTCCC GCGCAAGAAG ACGTGCCGT
751 CGTCCGGCTC AAATTTTGTG TCCTCGGCTT CTGCTCCCAG GCTGGACACT
801 GTCAGCGATG TGGAACCTGC ACAGAAGAAG GATGCGCTGA TTGTCAAAGA
851 AGCTCCAAA CCAAAGGCTC TTTCGGCCCC TGCAGCCCC GCTGTACAAG
901 AAGACCTTTG GGATTTCAAG AAATACATTG GTTTCGAGGA GCCCGTGGAG
951 GCCAAGGATG ATGGCTCGGC TGTGTCAGAT GATGCGGGTT CCTTTGAACA
1001 TCACCAGAA CATGATTCCG GACCTTTGGC AGGGGAGAAC GTCATGAACG
1051 TGGTCGTCGT TGCTGCTGAA TGTTCTCCCT GGTGCAAAAC AGGTGGTCTT
1101 GGAGATGTTG CGGGTGCTTT GCCCAAGGCT TTGGCTAAGA GAGGACATCG
1151 TGTATGGTT GTGGTACCAA GGTATGGGGA CTATGAGGAA GCCTACGATG
1201 TCGGAGTCCG AAAATACTAC AAGGCTGCTG GACAGGATAT GGAAGTGAAT
1251 TATTTCCATG CTTATATCGA TGGAGTGGAT TTTGTGTTCA TTGACGCTCC
1301 TCTCTCCGA CACCGTCAGC AAGACATTTA TGGGGGCAGC AGACAGGAAA
1351 TTATGAAGCG CATGATTTTG TTCTGCAAGG CCGCTGTCCA GGTTCCTTGG
1401 CACGTTCCAT GCGGCGGTGT CCTTACGGG GATGGAATC TGGTCTTCAT
1451 TGCAAATGAT TGGCACACGG CACTCCTGCC TGTCTATCTG AAAGCATATT
1501 ACAGGGACCA TGGTTTGATG CAATACAGTC GCTCCGTTAT GGTGATACAT
1551 AACATCGCTC ACCAGGGCCG TGGCCCTGTA GATGAATTCC CGTTCACCGA
1601 GTTGCTGAG CACTACCTGG AACACTTCAG ACTGTACGAC CCCGTGCGCG
1651 GTGAGCACGC CAACTACTTC GCCCGCGGCC TGAAGATGGC GGACCAAGTT
1701 GTCGTCGTGA GCCCCGGGTA CCTGTGGGAG CTGAAGACGG TGGAGGGCGG
1751 CTGGGGGCTT CACGACATCA TACGGCAGAA CGACTGGAAG ACCCGCGGCA
1801 TCGTGAACGG CATCGACAAC ATGGAGTGGA ACCCTGAGGT GGACGTCCAC
1851 CTGAAGTCGG ACGGCTACAC CAACTTCTCC CTGAAGACGC TGGACTCCGG
1901 CAAGCGGCAG TGCAAGGAGG CCCTGCAGCG CGAGCTGGGG CTGCAGGTCC
1951 GCGGCGACGT GCCGCTGCTC GGGTTCATCG GCGGCTGGA CCGGCAGAAG
2001 GGCGTGAGA TCATCGCGGA CGCGATGCCC TGGATCGTGA GCCAGGACGT
2051 GCAGCTGGTG ATGCTGGGCA CGGGGCGCCA CGACCTGGAG AGCATGCTGC
2101 AGCACTTCGA GCGGGAGCAC CACGACAAGG TGCGCGGGTG GGTGGGGTTC
2151 TCCGTGCGCC TGGCGCACCG GATCACGGCG GCGCGCGACG CGCTCCTCAT
2201 GCCCTCCCGG TTCGAGCCGT GCGGGCTGAA CCAGCTCTAC GCGATGGCCT
2251 ACGGCACCGT CCCCCTCGTG CACGCCGTCG GCGGCTTGAG GGATACCGTG
2301 CCGCCGTTCG ACCCTTCAA CCACTCCGGG CTCGGGTGGA CGTTCCACCG
2351 CGCCGAGGCG CACAAGCTGA TCGAGGCGCT CGGGCACTGC CTCCGCACCT
2401 ACCGGGACCA CAAGGAGAGC TGGAGGGGCC TCCAGGAGCG CCGCATGTCTG
2451 CAGGACTTCA GCTGGGAACA TGCCGCCAAG CTCTACGAGG ACGTCCCTCGT
2501 CCAGGCCAAG TACCAAGTGGT GAACGCTGCT ACCCGGTCCA GCGCCGCATG
2551 CGTGCATGAG AGGATGGAAA TCGCATTCG GCACTTGCAG ATTTGGCGCA
2601 CGCAGGAACG TGCCGTCCCTT CTTGATGAGA ACGCCGGCAT CCGCGAGGTT
2651 GAGACGCTGA TTCCGATCTG GTCCGTCGCA GAGTAGAGTG AAACGCTCCT
2701 TGTTGCAGGT ATATGGGAAT GTTTTTTTT CTTTTTTTTT GCGAGGGAGG
2751 TATATGGGAA TGTTAACTTG GTATTGTAAT GTGGTATGCT GTGTGCATTA
2801 TTACATCGGT TGTTGTTGCT TATTCTTGCT AGCTAAGTCG GAGGCCAAGA

```

10/23

2851 GCGAAAGCTA GCTCACATGT CTGATGTATG CAACTGACAT GGTTGGTTTG
2901 AAAAAAAAAA AAAAAAAAAA

11/23




FIGURE 10

12/23

Comparison of cDNA Sequences

	1				50
MK6827	-----GTG	CGTTTACCCC	ACACAGAGTA	CACTCCAACT	CCAGTCCAGT
MOREX	-----GTG	CGTTTACCCC	ACACAGAGTA	CACTCCAACT	CCAGTCCAGT
292	-----GTG	CGTTTACCCC	ACACAGAGTA	CACTCCAACT	CCAGTCCAGT
HIMALAYA	CCTCGAGGTG	CGTTTACCCC	ACACAGAGTA	CACTCCAACT	CCAGTCCAAAT
	51				100
MK6827	CCAGCCCACT	GCCGCTTCTG	CCCGCCCATC	GTACCGTCGC	CCGCCCCGAT
MOREX	CCAGCCCACT	GCCGCTTCTG	CCCGCCCATC	GTACCGTCGC	CCGCCCCGAT
292	CCAGCCCACT	GCCGCTTCTG	CCCGCCCATC	GTACCGTCGC	CCGCCCCGAT
HIMALAYA	CCAGCCCACT	GCCGCTTCTG	CCCGCCCATC	GTACCGTCGC	CCGCCCCGAT
	101	*** start codon			150
MK6827	CCCGGCCGCC	GCCATGTCGT	CGGCGGTCGC	GTCCCCCGCG	TCCTTCCTCG
MOREX	CCCGGCCGCC	GCCATGTCGT	CGGCGGTCGC	GTCCCCCGCG	TCCTTCCTCG
292	CCCGGCCGCC	GCCATGTCGT	CGGCGGTCGC	GTCCCCCGCG	TCCTTCCTCG
HIMALAYA	CCCGGCCGCC	GCCATGTCGT	CGGCGGTCGC	GTCCCCCGCG	TCCTTCCTCG
	151				200
MK6827	CGCTCGCGTC	CGCCTCGCCC	GGGAGATCAT	CACGGAGGAG	GGCGAGGGTG
MOREX	CGCTCGCGTC	CGCCTCGCCC	GGGAGATCAT	CACGGAGGAG	GGCGAGGGTG
292	CGCTCGCGTC	CGCCTCGCCC	GGGAGATCAT	CACGGAGGAG	GGCGAGGGTG
HIMALAYA	CGCTCGCGTC	CGCCTCGCCC	GGGAGATCAT	CACGGAGGAG	GGCGAGGGTG
	201			#	
MK6827	GGCGCGTCGC	CAACCCGCGC	TGGGGCCGGC	AGGCTGCAAT	GACGGCCGTC
MOREX	GGCGCGTCGC	CAACCCGCGC	TGGGGCCGGC	AGGCTGCAAT	GGCGGCCGTC
292	GGCGCGTCGC	CAACCCGCGC	TGGGGCCGGC	AGGCTGCAAT	GGCGGCCGTC
HIMALAYA	GGCGCGTCGC	CAACCCGCGC	TGGGGCCGGC	AGGCTGCAAT	GGCGGCCGTC
	251				300
MK6827	GCCGCTGCAG	CGCACGGGCTC	GCGACGGAGC	GGTGGCCGCG	CGCGCCGCCG
MOREX	GCCGCTGCAG	CGCACGGGCTC	GCGACGGAGC	GGTGGCCGCG	CGCGCCGCCG
292	GCCGCTGCAG	CGCACGGGCTC	GCGACGGAGC	GGTGGCCGCG	CGCGCCGCCG
HIMALAYA	GCCGCTGCAG	CGCACGGGCTC	GCGACGGAGC	GGTGGCCGCG	CGCGCCGCCG
	301				350
MK6827	GGATCGACGA	CGCCGCGCCC	GGTAGGCAGC	CCC GCGCTCG	CCGCTATGGC
MOREX	GGATCGACGA	CGCCGCGCCC	GGTAGGCAGC	CCC GCGCTCG	CCGCTATGGC
292	GGATCGACGA	CGCCGCGCCC	GGTAGGCAGC	CCC GCGCTCG	CCGCTATGGC
HIMALAYA	GGATCGACGA	CGCCGCGCCC	GGTAGGCAGC	CCC GCGCTCG	CCGCTATGGC
	351				400
MK6827	GCCGCCACCA	AGGTGCGGGA	TCCCGTCAAG	ACGCTCGATC	GCGACGCCGC
MOREX	GCCGCCACCA	AGGTGCGGGA	TCCCGTCAAG	ACGCTCGATC	GCGACGCCGC
292	GCCGCCACCA	AGGTGCGGGA	TCCCGTCAAG	ACGCTCGATC	GCGACGCCGC
HIMALAYA	GCCGCCACCA	AGGTGCGGGA	TCCCGTCAAG	ACGCTCGATC	GCGACGCCGC
	401				450
MK6827	GGAAGGTGGT	GGGCCGTCCC	CGCCGGCACC	GAGGCAGGAC	GCCGCCCGTC
MOREX	GGAAGGTGGT	GGGCCGTCCC	CGCCGGCACC	GAGGCAGGAC	GCCGCCCGTC
292	GGAAGGTGGT	GGGCCGTCCC	CGCCGGCACC	GAGGCAGGAC	GCCGCCCGTC
HIMALAYA	GGAAGGTGGT	GGGCCGTCCC	CGCCGGCACC	GAGGCAGGAC	GCCGCCCGTC
	451				500
MK6827	TGCCGAGTAA	GAACGGCACG	CTGATCAACG	GTGAGAACAA	ACCTACCGGC
MOREX	TGCCGAGTAA	GAACGGCACG	CTGATCAACG	GTGAGAACAA	ACCTACCGGC
292	TGCCGAGTAA	GAACGGCACG	CTGATCAACG	GTGAGAACAA	ACCTACCGGC
HIMALAYA	TGCCGAGTAA	GAACGGCACG	CTGATCAACG	GTGAGAACAA	ACCTACCGGC

FIGURE 11

13/23

	501				550
MK6827	GGCGGTGGCG	CGACTAAAGA	CAGCGGGCTG	CCCACACCCG	CACGCGCGCC
MOREX	GGCGGTGGCG	CGACTAAAGA	CAGCGGGCTG	CCCACACCCG	CACGCGCGCC
292	GGCGGTGGCG	CGACTAAAGA	CAGCGGGCTG	CCCACACCCG	CACGCGCGCC
HIMALAYA	GGCGGTGGCG	CGACTAAAGA	CAGCGGGCTT	GCCACACCCG	CACGCGCGCC
	551				600
MK6827	CCATCTGTCA	ATCCAGAACA	GAGTACCGGT	GAACGGTGAA	AACAAACATA
MOREX	CCATCTGTCA	ATCCAGAACA	GAGTACCGGT	GAACGGTGAA	AACAAACATA
292	CCATCTGTCA	ATCCAGAACA	GAGTACCGGT	GAACGGTGAA	AACAAACATA
HIMALAYA	CCATCTGTCA	ATCCAAAACA	GAGTACCGGT	GAACGGTGAA	AACAAACATA
	601				650
MK6827	AGGTCGCCTC	GCCGCCGACC	AGCATAGTGG	ATGTCGCGTC	TCCGGGTTC
MOREX	AGGTCGCCTC	GCCGCCGACC	AGCATAGTGG	ATGTCGCGTC	TCCGGGTTC
292	AGGTCGCCTC	GCCGCCGACC	AGCATAGTGG	ATGTCGCGTC	TCCGGGTTC
HIMALAYA	AGGTCGCCTC	GCCGCCGACC	AGCATAGTGG	ATGTCGCGTC	TCCGGGTTC
	651				700
MK6827	GCAGCTAACA	TTTCCATCAG	TAACAAGGTG	CCGCCGTCCG	TTGTCCCAGC
MOREX	GCAGCTAACA	TTTCCATCAG	TAACAAGGTG	CCGCCGTCCG	TTGTCCCAGC
292	GCAGCCAACA	TTTCCATCAG	TAACAAGGTG	CCGCCGTCCG	TTGTCCCAGC
HIMALAYA	GCAGCTAACA	TTTCCATCAG	TAACAAGGTG	CCGCCGTCCG	TTGTCCCAGC
	701				750
MK6827	CAAGAAGACG	CCGCCGTCTG	CCGTTTTCCC	GGCCAAGAAG	GCGCCGCCGT
MOREX	CAAGAAGACG	CCGCCGTCTG	CCGTTTTCCC	GGCCAAGAAG	GCGCCGCCGT
292	CAAGAAGACG	CCGCCGTCTG	CCGTTTTCCC	GGCCAAGAAG	GCGCCGCCGT
HIMALAYA	CAAGAAGACG	CCGCCGTCTG	CCGTTTTCCC	GGCCAAGAAG
	751				800
MK6827	CGTCCGTTGT	CCCGGCCAAG	AAGACGCTGC	CGTCGTCCGG	CTCAAATTTT
MOREX	CGTCCGTTGT	CCCGGCCAAG	AAGACGCTGC	CGTCGTCCGG	CTCAAATTTT
292	CGTCCGTTGT	CCCGGCCAAG	AAGACGCTGC	CGTCGTCCGG	CTCAAATTTT
HIMALAYAACGCTGC	CGTCGTCCGG	CTCAAATTTT
	801				850
MK6827	GTGTCCTCGG	CCTCTGCTCC	CAGGCTGGAC	ACTGTCAGCG	ATGTGGAAC
MOREX	GTGTCCTCGG	CCTCTGCTCC	CAGGCTGGAC	ACTGTCAGCG	ATGTGGAAC
292	GTGTCCTCGG	CCTCTGCTCC	CAGGCTGGAC	ACTGTCAGCG	ATGTGGAAC
HIMALAYA	GTGTCCTCGG	CCTCTGCTCC	CAGGCTGGAC	ACTGTCAGCG	ATGTGGAAC
	851				900
MK6827	TGCACAGAAG	AAGGATGCGC	TGATTGTCAA	AGAAGCTCCA	AAACCAAAGG
MOREX	TGCACAGAAG	AAGGATGCGC	TGATTGTCAA	AGAAGCTCCA	AAACCAAAGG
292	TGCACAGAAG	AAGGATGCGC	TGATTGTCAA	AGAAGCTCCA	AAACCAAAGG
HIMALAYA	TGCACAGAAG	AAGGATGCGC	TGATTGTCAA	AGAAGCTCCA	AAACCAAAGG
	901				950
MK6827	CTCTTTTCGGC	CCCTGCAGCC	CCCGCTGTAC	AAGAAGACCT	TTGGGATTTT
MOREX	CTCTTTTCGGC	CCCTGCAGCC	CCCGCTGTAC	AAGAAGACCT	TTGGGATTTT
292	CTCTTTTCGGC	CCCTGCAGCC	CCCGCTGTAC	AAGAAGACCT	TTGGGATTTT
HIMALAYA	CTCTTTTCGGC	CCCTGCAGCC	CCCGCTGTAC	AAGAAGACCT	TTGGGATTTT
	951				1000
MK6827	AAGAAATACA	TTGGTTTCGA	GGAGCCCGTG	GAGGCCAAGG	ATGATGGCTC
MOREX	AAGAAATACA	TTGGTTTCGA	GGAGCCCGTG	GAGGCCAAGG	ATGATGGCTC
292	AAGAAATACA	TTGGTTTCGA	GGAGCCCGTG	GAGGCCAAGG	ATGATGGCTC
HIMALAYA	AAGAAATACA	TTGGTTTCGA	GGAGCCCGTG	GAGGCCAAGG	ATGATGGCTC

14/23

	1001		1050
MK6827	GGCTGTTGCA GATGATGCGG GTTCCTTTGA ACATCACCAG AATCATGATT		
MOREX	GGCTGTTGCA GATGATGCGG GTTCCTTTGA ACATCACCAG AATCATGATT		
292	GGCTGTTGCA GATGATGCGG GTTCCTTTGA ACATCACCAG AATCATGATT		
HIMALAYA	GGCTGTTGCA GATGATGCGG GTTCCTTTGA ACATCACCAG AATCATGATT		
	1051		1100
MK6827	CCGGACCTTT GGCAGGGGAG AACGTCATGA ACGTGGTCGT CGTTGCTGCT		
MOREX	CCGGACCTTT GGCAGGGGAG AACGTCATGA ACGTGGTCGT CGTTGCTGCT		
292	CCGGACCTTT GGCAGGGGAG AACGTCATGA ACGTGGTCGT CGTTGCTGCT		
HIMALAYA	CCGGACCTTT GGCAGGGGAG AACGTCATGA ACGTGGTCGT CGTTGCTGCT		
	1101		1150
MK6827	GAATGTTCTC CCTGGTGCAA AACAGGTGGT CTTGGAGATA TTGCGGGTGC		
MOREX	GAATGTTCTC CCTGGTGCAA AACAGGTGGT CTTGGAGATG TTGCGGGTGC		
292	GAATGTTCTC CCTGGTGCAA AACAGGTGGT CTTGGAGATG TTGCGGGTGC		
HIMALAYA	GAATGTTCTC CCTGGTGCAA AACAGGTGGT CTTGGAGATG TTGCGGGTGC		
	1151		1200
MK6827	TTTGCCCAAG GCTTTGGCTA AGAGAGGACA TCGTGTTATG GTTGTGGTAC		
MOREX	TTTGCCCAAG GCTTTGGCTA AGAGAGGACA TCGTGTTATG GTTGTGGTAC		
292	TTTGCCCAAG GCTTTGGCTA AGAGAGGACA TCGTGTTATG GTTGTGGTAC		
HIMALAYA	TTTGCCCAAG GCTTTGGCTA AGAGAGGACA TCGTGTTATG GTTGTGGTAC		
	1201		1250
MK6827	CAAGGTATGG GGAATATGAG GAAGCCTACG ATGTCGGAGT CCGAAAATAC		
MOREX	CAAGGTATGG GGAATATGAG GAAGCCTACG ATGTCGGAGT CCGAAAATAC		
292	CAAGGTATGG GGAATATGAG GAAGCCTACG ATGTCGGAGT CCGAAAATAC		
HIMALAYA	CAAGGTATGG GGAATATGAG GAAGCCTACG ATGTCGGAGT CCGAAAATAC		
	1251		1300
MK6827	TACAAGGCTG CTGGACAGGA TATGGAAGTG AATTATTTCC ATGCTTATAT		
MOREX	TACAAGGCTG CTGGACAGGA TATGGAAGTG AATTATTTCC ATGCTTATAT		
292	TACAAGGCTG CTGGACAGGA TATGGAAGTG AATTATTTCC ATGCTTATAT		
HIMALAYA	TACAAGGCTG CTGGACAGGA TATGGAAGTG AATTATTTCC ATGCTTATAT		
	1301		1350
MK6827	CGATGGAGTG GATTTTGTGT TCATTGACGC TCCTCTCTTC CGACACCGTC		
MOREX	CGATGGAGTG GATTTTGTGT TCATTGACGC TCCTCTCTTC CGACACCGTC		
292	CGATGGAGTG GATTTTGTGT TCATTGACGC TCCTCTCTTC CGACACCGTC		
HIMALAYA	CGATGGAGTG GATTTTGTGT TCATTGACGC TCCTCTCTTC CGACACCGTC		
	1351		1400
MK6827	AGCAAGACAT TTATGGGGGC AGCAGACAGG AAATTATGAA GCGCATGATT		
MOREX	AGCAAGACAT TTATGGGGGC AGCAGACAGG AAATTATGAA GCGCATGATT		
292	AGCAAGACAT TTATGGGGGC AGCAGACAGG AAATTATGAA GCGCATGATT		
HIMALAYA	AGCAAGACAT TTATGGGGGC AGCAGACAGG AAATTATGAA GCGCATGATT		
	1401		1450
MK6827	TTGTTCTGCA AGGCCGCTGT CGAGGTTTCTT TGGCACGTTT CATGCGGCGG		
MOREX	TTGTTCTGCA AGGCCGCTGT CGAGGTTTCTT TGGCACGTTT CATGCGGCGG		
292	TTGTTCTGCA AGGCCGCTGT CGAGGTTTCTT TGGCACGTTT CATGCGGCGG		
HIMALAYA	TTGTTCTGCA AGGCCGCTGT CGAGGTTTCTT TGGCACGTTT CATGCGGCGG		
	1451		1500
MK6827	TGTCCCTTAC GGGGATGGAA ATCTGGTCTT CATTGCAAAT GATTGGCACA		
MOREX	TGTCCCTTAC GGGGATGGAA ATCTGGTCTT CATTGCAAAT GATTGGCACA		
292	TGTCCCTTAC GGGGATGGAA ATCTGGTCTT CATTGCAAAT GATTGGCACA		
HIMALAYA	TGTCCCTTAC GGGGATGGAA ATCTGGTCTT CATTGCAAAT GATTGGCACA		

15/23

	1501		1550
MK6827	CGGCACTCCT GCCTGTCTAT CTGAAAGCAT ATTACAGGGA CCATGGTTTG		
MOREX	CGGCACTCCT GCCTGTCTAT CTGAAAGCAT ATTACAGGGA CCATGGTTTG		
292	CGGCACTCCT GCCTGTCTAT CTGAAAGCAT ATTACAGGGA CCATGGTTTG		
HIMALAYA	CGGCACTCCT GCCTGTCTAT CTGAAAGCAT ATTACAGGGA CCATGGTTTG		
	1551		1600
MK6827	ATGCAATACA GTCGCTCCGT TATGGTGATA CATAACATCG CTCACCAGGG		
MOREX	ATGCAATACA GTCGCTCCGT TATGGTGATA CATAACATCG CTCACCAGGG		
292	ATGCAATACA GTCGCTCCGT TATGGTGATA CATAACATCG CTCACCAGGG		
HIMALAYA	ATGCAATACA GTCGCTCCGT TATGGTGATA CATAACATCG CTCACCAGGG		
	1601		1650
MK6827	CCGTGGCCCT GTAGATGAAT TCCCGTTCAC CGAGTTGCCT GAGCACTACC		
MOREX	CCGTGGCCCT GTAGATGAAT TCCCGTTCAC CGAGTTGCCT GAGCACTACC		
292	CCGTGGCCCT GTAGATGAAT TCCCGTTCAC CGAGTTGCCT GAGCACTACC		
HIMALAYA	CCGTGGCCCT GTAGATGAAT TCCCGTTCAC CGAGTTGCCT GAGCACTACC		
	1651		1700
MK6827	TGGAACACTT CAGACTGTAC GACCCCGTCG GCGGTGAGCA CGCCAACTAC		
MOREX	TGGAACACTT CAGACTGTAC GACCCCGTCG GCGGTGAGCA CGCCAACTAC		
292	TGGAACACTT CAGACTGTAC GACCCCGTCG GCGGTGAGCA CGCCAACTAC		
HIMALAYA	TGGAACACTT CAGACTGTAC GACCCCGTCG GCGGTGAGCA CGCCAACTAC		
	1701		1750
MK6827	TTCGCCGCCG GCCTGAAGAT GGCGGACCAG GTTGTCTGTCG TGAGCCCCCGG		
MOREX	TTCGCCGCCG GCCTGAAGAT GGCGGACCAG GTTGTCTGTCG TGAGCCCCCGG		
292	TTCGCCGCCG GCCTGAAGAT GGCGGACCAG GTTGTCTGTCG TGAGCCCCCGG		
HIMALAYA	TTCGCCGCCG GCCTGAAGAT GGCGGACCAG GTTGTCTGTCG TGAGCCCCCGG		
	1751		1800
MK6827	GTACCTGTGG GAGCTGAAGA CGGTGGAGGG CGGCTGGGGG CTTACGACA		
MOREX	GTACCTGTGG GAGCTGAAGA CGGTGGAGGG CGGCTGGGGG CTTACGACA		
292	GTACCTGTGG GAGCTGAAGA CGGTGGAGGG CGGCTGGGGG CTTACGACA		
HIMALAYA	GTACCTGTGG GAGCTGAAGA CGGTGGAGGG CGGCTGGGGG CTTACGACA		
	1801		1850
MK6827	TCATACGGCA GAACGACTGG AAGACCCGCG GCATCGTGAA CGGCATCGAC		
MOREX	TCATACGGCA GAACGACTGG AAGACCCGCG GCATCGTGAA CGGCATCGAC		
292	TCATACGGCA GAACGACTGG AAGACCCGCG GCATCGTGAA CGGCATCGAC		
HIMALAYA	TCATACGGCA GAACGACTGG AAGACCCGCG GCATCGTGAA CGGCATCGAC		
	1851	&	1900
MK6827	AACATGGAGT GGAACCCCTGA GGTGGACGTC CACCTGAAGT CGGACGGCTA		
MOREX	AACATGGAGT GGAACCCCTGA GGTGGACGTC CACCTGAAGT CGGACGGCTA		
292	AACATGGAGT GGAACCCCTGA GGTGGACGTC CACCTGAAGT CGGACGGCTA		
HIMALAYA	AACATGGAGT GGAACCCCTGA GGTGGACGTC CACCTGAAGT CGGACGGCTA		
	1901		1950
MK6827	CACCAACTTC TCCCTGAAGA CGCTGGACTC CGGCAAGCGG CAGTGCAAGG		
MOREX	CACCAACTTC TCCCTGAAGA CGCTGGACTC CGGCAAGCGG CAGTGCAAGG		
292	CACCAACTTC TCCCTGAAGA CGCTGGACTC CGGCAAGCGG CAGTGCAAGG		
HIMALAYA	CACCAACTTC TCCCTGAAGA CGCTGGACTC CGGCAAGCGG CAGTGCAAGG		
	1951		2000
MK6827	AGGCCCTGCA GCGCGAGCTG GGGCTGCAGG TCCGCGGCGA CGTGCCGCTG		
MOREX	AGGCCCTGCA GCGCGAGCTG GGGCTGCAGG TCCGCGGCGA CGTGCCGCTG		
292	AGGCCCTGCA GCGCGAGCTG GGGCTGCAGG TCCGCGGCGA CGTGCCGCTG		
HIMALAYA	AGGCCCTGCA GCGCGAGCTG GGGCTGCAGG TCCGCGGCGA CGTGCCGCTG		

16/23

	2001				2050
MK6827	CTCGGGTTCA	TCGGGCGGCT	GGACGGGCAG	AAGGGCGTGG	AGATCATCGC
MOREX	CTCGGGTTCA	TCGGGCGGCT	GGACGGGCAG	AAGGGCGTGG	AGATCATCGC
292	CTCGGGTTCA	TCGGGCGGCT	GGACGGGCAG	AAGGGCGTGG	AGATCATCGC
HIMALAYA	CTCGGGTTCA	TCGGGCGGCT	GGACGGGCAG	AAGGGCGTGG	AGATCATCGC
	2051				2100
MK6827	GGACGCGATG	CCCTGGATCG	TGAGCCAGGA	CGTGCAGCTG	GTGATGCTGG
MOREX	GGACGCGATG	CCCTGGATCG	TGAGCCAGGA	CGTGCAGCTG	GTGATGCTGG
292	GGACGCGATG	CCCTGGATCG	TGAGCCAGGA	CGTGCAGCTG	GTGATGCTGG
HIMALAYA	GGACGCGATG	CCCTGGATCG	TGAGCCAGGA	CGTGCAGCTG	GTGATGCTGG
	2101				2150
MK6827	GCACGGGGCG	CCACGACCTG	GAGAGCATGC	TGCAGCACTT	CGAGCGGGAG
MOREX	GCACGGGGCG	CCACGACCTG	GAGAGCATGC	TGCAGCACTT	CGAGCGGGAG
292	GCACGGGGCG	CCACGACCTG	GAGAGCATGC	TGCAGCACTT	CGAGCGGGAG
HIMALAYA	GCACGGGGCG	CCACGACCTG	GAGAGCATGC	TGCAGCACTT	CGAGCGGGAG
	2151				2200
MK6827	CACCACGACA	AGGTGCGCGG	GTGGGTGGGG	TTCTCCGTGC	GCCTGGCGCA
MOREX	CACCACGACA	AGGTGCGCGG	GTGGGTGGGG	TTCTCCGTGC	GCCTGGCGCA
292	CACCACGACA	AGGTGCGCGG	GTGGGTGGGG	TTCTCCGTGC	GCCTGGCGCA
HIMALAYA	CACCACGACA	AGGTGCGCGG	GTGGGTGGGG	TTCTCCGTGC	GCCTGGCGCA
	2201				2250
MK6827	CCGGATCACG	GCGGGCGCCG	ACGCGCTCCT	CATGCCCTCC	CGGTTCCGAGC
MOREX	CCGGATCACG	GCGGGCGCCG	ACGCGCTCCT	CATGCCCTCC	CGGTTCCGAGC
292	CCGGATCACG	GCGGGCGCCG	ACGCGCTCCT	CATGCCCTCC	CGGTTCCGAGC
HIMALAYA	CCGGATCACG	GCGGGCGCCG	ACGCGCTCCT	CATGCCCTCC	CGGTTCCGAGC
	2251				2300
MK6827	CGTGCGGGCT	GAACCAGCTC	TACGCGATGG	CCTACGGCAC	CATCCCTGTC
MOREX	CGTGCGGGCT	GAACCAGCTC	TACGCGATGG	CCTACGGCAC	CATCCCTGTC
292	CGTGCGGGCT	GAACCAGCTC	TACGCGATGG	CCTACGGCAC	CATCCCTGTC
HIMALAYA	CGTGCGGGCT	GAACCAGCTC	TACGCGATGG	CCTACGGCAC	CGTCCCCGTC
	2301				2350
MK6827	GTGCACGCCG	TCGGCGGCCT	GAGGGATAACC	GTGCCGCCGT	TCGACCCCTT
MOREX	GTGCACGCCG	TCGGCGGCCT	GAGGGATAACC	GTGCCGCCGT	TCGACCCCTT
292	GTGCACGCCG	TCGGCGGCCT	GAGGGATAACC	GTGCCGCCGT	TCGACCCCTT
HIMALAYA	GTGCACGCCG	TCGGCGGCCT	GAGGGATAACC	GTGCCGCCGT	TCGACCCCTT
	2351				2400
MK6827	CAACCACTCC	GGGCTCGGGT	GGACGTTCTGA	CCGCGCCGAG	GCGCACAAGC
MOREX	CAACCACTCC	GGGCTCGGGT	GGACGTTCTGA	CCGCGCCGAG	GCGCACAAGC
292	CAACCACTCC	GGGCTCGGGT	GGACGTTCTGA	CCGCGCCGAG	GCGCACAAGC
HIMALAYA	CAACCACTCC	GGGCTCGGGT	GGACGTTCTGA	CCGCGCCGAG	GCGCACAAGC
	2401				2450
MK6827	TGATCGAGGC	GCTCGGGCAC	TGCCTCCGCA	CCTACCGGGA	CCACAAGGAG
MOREX	TGATCGAGGC	GCTCGGGCAC	TGCCTCCGCA	CCTACCGGGA	CCACAAGGAG
292	TGATCGAGGC	GCTCGGGCAC	TGCCTCCGCA	CCTACCGGGA	CCACAAGGAG
HIMALAYA	TGATCGAGGC	GCTCGGGCAC	TGCCTCCGCA	CCTACCGGGA	CCACAAGGAG
	2451				2500
MK6827	AGCTGGAGGG	GCCTCCAGGA	GCGCGGCATG	TCGCAGGACT	TCAGCTGGGA
MOREX	AGCTGGAGGG	GCCTCCAGGA	GCGCGGCATG	TCGCAGGACT	TCAGCTGGGA
292	AGCTGGAGGG	GCCTCCAGGA	GCGCGGCATG	TCGCAGGACT	TCAGCTGGGA
HIMALAYA	AGCTGGAGGG	GCCTCCAGGA	GCGCGGCATG	TCGCAGGACT	TCAGCTGGGA

17/23

	2501		2550
MK6827	ACATGCCGCC AAGCTCTACG AGGACGTCCT CGTCCAGGCC AAGTACCACT		
MOREX	ACATGCCGCC AAGCTCTACG AGGACGTCCT CGTCCAGGCC AAGTACCACT		
292	ACATGCCGCC AAGCTCTACG AGGACGTCCT CGTCCAGGCC AAGTACCACT		
HIMALAYA	ACATGCCGCC AAGCTCTACG AGGACGTCCT CGTCCAGGCC AAGTACCACT		
	2551		2600
	*** stop codon		
MK6827	GGTGAACGCT GCTACCCGGT CCAGCCCCGC ATGCGTGCAT GAGAGGATGG		
MOREX	GGTGAACGCT GCTACCCGGT CCAGCCCCGC ATGCGTGCAT GAGAGGATGG		
292	GGTGAACGCT GCTACCCGGT CCAGCCCCGC ATGCGTGCAT GAGAGGATGG		
HIMALAYA	GGTGAACGCT GCTACCCGGT CCAGCCCCGC ATGCGTGCAT GAGAGGATGG		
	2601		2650
MK6827	AAATGCGCAT TGCGCACCTG CAGATTGCGC GCATGCAGGA ACGTGCCGTC		
MOREX	AAATGCGCAT TGCGCACCTG CAGATTGCGC GCATGCAGGA ACGTGCCGTC		
292	AAATGCGCAT TGCGCACCTG CAGATTGCGC GCATGCAGGA ACGTGCCGTC		
HIMALAYA	AAATGCGCAT TGCGCACCTG CAGATTGCGC GCATGCAGGA ACGTGCCGTC		
	2651		2700
MK6827	CTTCTTGATG GGAACGCCGG CATCCGCGAG GTTGAGACGC TGATTCCGAT		
MOREX	CTTCTTGATG GGAACGCCGG CATCCGCGAG GTTGAGACGC TGATTCCGAT		
292	CTTCTTGATG GGAACGCCGG CATCCGCGAG GTTGAGACGC TGATTCCGAT		
HIMALAYA	CTTCTTGATG GGAACGCCGG CATCCGCGAG GTTGAGACGC TGATTCCGAT		
	2701		2750
MK6827	CTGGTCCGTC GCAGAGTAGA GTGAAACGCT CCTTGTTGCA GGTATATGGG		
MOREX	CTGGTCCGTC GCAGAGTAGA GTGAAACGCT CCTTGTTGCA GGTATATGGG		
292	CTGGTCCGTC GCAGAGTAGA GTGAAACGCT CCTTGTTGCA GGTATATGGG		
HIMALAYA	CTGGTCCGTC GCAGAGTAGA GTGAAACGCT CCTTGTTGCA GGTATATGGG		
	2751		2800
MK6827	AATGTTTTTT TTTTCC.TTT TTTTTTTTGC GAGGGAGGTA TATGGGAATG		
MOREX	AATGTTTTTT TTTTCC.TTT TTTTTTTTGC GAGGGAGGTA TATGGGAATG		
292	AATGTTTTTT TT..CC...T TTTTTTTTGC GAGGGAGGTA TATGGGAATG		
HIMALAYA	AATGTTTTTT TT..CC...T TTTTTTTTGC GAGGGAGGTA TATGGGAATG		
	2801		2850
MK6827	TTAACTTGGT ATTGTAATGT GGTATGCTGT GTGCATTATT ACATCGGTTG		
MOREX	TTAACTTGGT ATTGTAATGT GGTATGCTGT GTGCATTATT ACATCGGTTG		
292	TTAACTTGGT ATTGTAATGT GGTATGCTGT GTGCATTATT ACATCGGTTG		
HIMALAYA	TTAACTTGGT ATTGTAATGT GGTATGCTGT GTGCATTATT ACATCGGTTG		
	2851		2900
MK6827	TTGTTGCTTA TTCTTGCTAG CTAAGTCGGA GGCCAAGAGC GAAAGCTAGC		
MOREX	TTGTTGCTTA TTCTTGCTAG CTAAGTCGGA GGCCAAGAGC GAAAGCTAGC		
292	TTGTTGCTTA TTCTTGCTAG CTAAGTCGGA GGCCAAGAGC GAAAGCTAGC		
HIMALAYA	TTGTTGCTTA TTCTTGCTAG CTAAGTCGGA GGCCAAGAGC GAAAGCTAGC		
	2901		2950
MK6827	TCACATGTCT GATGTATGCA AGTGACATGG TTGGTTTGGT TGTGCAGTGC		
MOREX	TCACATGTCT GATGTATGCA AGTGACATGG TTGGTTTGGT TGTGCAGTGC		
292	TCACATGTCT GATGTATGCA AGTGACATGG TTGGTTTGGT TGTGCAGTGC		
HIMALAYA	TCACATGTCT GATGTATGCA AGTGACATGG TTGGTTTGGT TGTGCAGTGC		
	2951		
MK6827	AAACGGCA		
MOREX	AAACGGCA		
292	AAACGGCA		
HIMALAYA	AAAAA		

18/23

Comparison of SSII Amino Acid Sequences

Morex Himalaya 292 MK6827	1	MK6827 mutation #			
		MSSAVASPAS	FLALASASPG	RSSRRRARVG	ASPTRAGAGR LQWRPSPLQR
		MSSAVASPAS	FLALASASPG	RSSRRRARVG	ASPTRAGAGR LQWRPSPLQR
		MSSAVASPAS	FLALASASPG	RSSRRRARVG	ASPTRAGAGR LQWRPSPLQR
Morex Himalaya 292 MK6827	51				
		TARDGAVAAR	AAGIDDAAPG	RQPRARRYGA	ATKVADPVKT LDRDAAEGGG
		TARDGAVAAR	AAGIDDAAPG	RQPRARRYGA	ATKVADPVKT LDRDAAEGGG
		TARDGAVAAR	AAGIDDAAPG	RQPRARRYGA	ATKVADPVKT LDRDAAEGGG
Morex Himalaya 292 MK6827	101				
		PSPPAPRQDA	ARLPSKNGTL	INGENKPTGG	GGATKDSGLP TPARAPHLST
		PSPPAPRQDA	ARLPSKNGTL	INGENKPTGG	GGATKDSGLP TPARAPHLST
		PSPPAPRQDA	ARLPSKNGTL	INGENKPTGG	GGATKDSGLP TPARAPHLST
Morex Himalaya 292 MK6827	151				
		QNRVPVNGEN	KHKVASPPTS	IVDVASPGSA	ANISISNKVP PSVVPAKKTTP
		QNRVPVNGEN	KHKVASPPTS	IVDVASPGSA	ANISISNKVP PSVVPAKKTTP
		QNRVPVNGEN	KHKVASPPTS	IVDVASPGSA	ANISISNKVP PSVVPAKKTTP
Morex Himalaya 292 MK6827	201				
		PSSVFPKKT	LPSSGSNFVS	SASAPRLDTV	SDVELAQKKD ALIVKEAPKP
		PSSVFPKKT	LPSSGSNFVS	SASAPRLDTV	SDVELAQKKD ALIVKEAPKP
		PSSVFPKKT	LPSSGSNFVS	SASAPRLDTV	SDVELAQKKD ALIVKEAPKP
Morex Himalaya 292 MK6827	251				
		KALSAPAAPA	VQEDLWDFKK	YIGFEEPVEA	KDDGSAVADD AGSFEHHQNH
		KALSAPAAPA	VQEDLWDFKK	YIGFEEPVEA	KDDGSAVADD AGSFEHHQNH
		KALSAPAAPA	VQEDLWDFKK	YIGFEEPVEA	KDDGSAVADD AGSFEHHQNH
Morex Himalaya 292 MK6827	301				
		DSGPLAGENV	MNVVVVAEEC	SPWCKTGGLG	DVAGALPKAL AKRGHRVMV
		DSGPLAGENV	MNVVVVAEEC	SPWCKTGGLG	DVAGALPKAL AKRGHRVMV
		DSGPLAGENV	MNVVVVAEEC	SPWCKTGGLG	DVAGALPKAL AKRGHRVMV
Morex Himalaya 292 MK6827	351				
		VPRYGDYEEA	YDVGVRKYYK	AAGQDMEVNY	FHAYIDGVDF VFIDAPLFRH
		VPRYGDYEEA	YDVGVRKYYK	AAGQDMEVNY	FHAYIDGVDF VFIDAPLFRH
		VPRYGDYEEA	YDVGVRKYYK	AAGQDMEVNY	FHAYIDGVDF VFIDAPLFRH
Morex Himalaya 292 MK6827	401				
		RQDDIYGGS	QEIMKRMILF	CKAAVEVPWH	VPCGGVPYGD GNLVFIANDW
		RQDDIYGGS	QEIMKRMILF	CKAAVEVPWH	VPCGGVPYGD GNLVFIANDW
		RQDDIYGGS	QEIMKRMILF	CKAAVEVPWH	VPCGGVPYGD GNLVFIANDW
Morex Himalaya 292 MK6827	451				
		HTALLPVYLK	AYYRDHGLMQ	YSRSVMVIHN	IAHQGRGPVD EFPFTELPEH
		HTALLPVYLK	AYYRDHGLMQ	YSRSVMVIHN	IAHQGRGPVD EFPFTELPEH
		HTALLPVYLK	AYYRDHGLMQ	YSRSVMVIHN	IAHQGRGPVD EFPFTELPEH
Morex Himalaya 292 MK6827	501				
		YLEHFRLYDP	VGGEHANYFA	AGLKMAQV	VVSPGYLWEL KTVEGGWGLH
		YLEHFRLYDP	VGGEHANYFA	AGLKMAQV	VVSPGYLWEL KTVEGGWGLH
		YLEHFRLYDP	VGGEHANYFA	AGLKMAQV	VVSPGYLWEL KTVEGGWGLH

FIGURE 12

19/23

	551		\$ 292 mutation		600
Morex	DIIRQNDWKT	RGIVNGIDNM	EWNPEVDVHL	KSDGYTNFSL	KTLD SGKRQC
Himalaya	DIIRQNDWKT	RGIVNGIDNM	EWNPEVDVHL	KSDGYTNFSL	KTLD SGKRQC
292	DIIRQNDWKT	RGIVNGIDNM	E*NPEVDVHL	KSDGYTNFSL	KTLD SGKRQC
MK6827	DIIRQNDWKT	RGIVNGIDNM	EWNPEVDVHL	KSDGYTNFSL	KTLD SGKRQC
	601				650
Morex	KEALQRELGL	QVRGDVPLLG	FIGRLDGQKG	VEIIADAMPW	IVSQDVQLVM
Himalaya	KEALQRELGL	QVRGDVPLLG	FIGRLDGQKG	VEIIADAMPW	IVSQDVQLVM
292	KEALQRELGL	QVRGDVPLLG	FIGRLDGQKG	VEIIADAMPW	IVSQDVQLVM
MK6827	KEALQRELGL	QVRGDVPLLG	FIGRLDGQKG	VEIIADAMPW	IVSQDVQLVM
	651				700
Morex	LGTGRHDLES	MLQHFEREHH	DKVRGWVGFS	VRLAHRITAG	ADALLMPSRF
Himalaya	LGTGRHDLES	MLQHFEREHH	DKVRGWVGFS	VRLAHRITAG	ADALLMPSRF
292	LGTGRHDLES	MLQHFEREHH	DKVRGWVGFS	VRLAHRITAG	ADALLMPSRF
MK6827	LGTGRHDLES	MLQHFEREHH	DKVRGWVGFS	VRLAHRITAG	ADALLMPSRF
	701				750
Morex	EPCGLNQLYA	MAYGTIPVVH	AVGGLRDTVP	PFDPFNHSG	GWTFDRAEAAH
Himalaya	EPCGLNQLYA	MAYGTIPVVH	AVGGLRDTVP	PFDPFNHSG	GWTFDRAEAAH
292	EPCGLNQLYA	MAYGTIPVVH	AVGGLRDTVP	PFDPFNHSG	GWTFDRAEAAH
MK6827	EPCGLNQLYA	MAYGTIPVVH	AVGGLRDTVP	PFDPFNHSG	GWTFDRAEAAH
	751				800
Morex	KLIEALGHCL	RTYRDHKESW	RGLQERGMSQ	DFSWEHAACL	YEDVLVQAKY
Himalaya	KLIEALGHCL	RTYRDHKESW	RGLQERGMSQ	DFSWEHAACL	YEDVLVQAKY
292	KLIEALGHCL	RTYRDHKESW	RGLQERGMSQ	DFSWEHAACL	YEDVLVQAKY
MK6827	KLIEALGHCL	RTYRDHKESW	RGLQERGMSQ	DFSWEHAACL	YEDVLVQAKY
	801				
Morex	QW*				
Himalaya	QW*				
292	QW*				
MK6827	QW*				

20/23


Position 242 of barley SSII
cDNA sequence

Himalaya

.....CGGCAGGCTGCAATGGCGGCCGTCGCCGCT.....

MK6827

.....CGGCAGGCTGCAATGACGGGCCGTCGCCGCT.....



Position 1829 of barley SSII
cDNA sequence

292

.....GACAACATGGAGTGAAACCCTGAGGTGGACGTCCA.....

Himalaya

.....GACAACATGGAGTGGAACCCTGAGGTGGACGTCCA.....

FIGURE 13






FIGURE 14



22/23




FIGURE 15

23/23




FIGURE 16

INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU01/01452

A. CLASSIFICATION OF SUBJECT MATTERInt. Cl. ⁷: A01H 5/00, 5/10; C12N 15/29; C08L 3/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC (WPIDS) AND CHEMICAL ABSTRACTS - KEYWORDS BELOW

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
SEE BELOWElectronic data base consulted during the international search (name of data base and, where practicable, search terms used)
GENBANK, EMBL, WPIDS, CA, MEDLINE, AGRICOLA. Keywords: starch, synthase, II, 2, synthaseII, synthase2, ssII, ss2, ss, barley, hordeum, wheat, rice, maize, high()amylose, resistant()starch**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	National Plant Germplasm System (http://www.ars-grin.gov/npgs/), GRIN System accession no. GSHO 2476, 23 June 1997	1-139,143
X	WO A 00/66745 (COMMONWEALTH SCIENTIFIC AND INDUSTRIAL RESEARCH ORGANISATION ET AL.) 9 November 2000	1-143
X	EMBL abstract accession no. AF155217, Li et al., 7 September 1999, "Triticum aestivum starch synthase IIA mRNA, complete cds."	140-141

☒ Further documents are listed in the continuation of Box C ☒ See patent family annex

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search.

4 January 2002

Date of mailing of the international search report

21 JAN 2002

Name and mailing address of the ISA/AU

AUSTRALIAN PATENT OFFICE
PO BOX 200, WODEN ACT 2606, AUSTRALIA
E-mail address: pct@ipaustalia.gov.au
Facsimile No (02) 6285 3929

Authorized officer

Chris Luton

Telephone No : (02) 6283 2256

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/AU01/01452

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report		Patent Family Member	
WO	200066745	AU	200040924
			END OF ANNEX